



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

### **Usage guidelines**


Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.


### **About Google Book Search**

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>










# EDUCATIONAL MATHEMATICAL WORKS.

By **I. TODHUNTER, M.A.**

*Fellow and Assistant Tutor of St John's College, Cambridge.*



1. **A Treatise on Algebra,**  
For the use of Students in the Universities and in Schools.  
Crown 8vo. cloth, 7s. 6d.
2. **A Treatise on Plane Co-ordinate Geometry,**  
As applied to the Straight Line and the CONIC SECTIONS.  
With numerous Examples. Second Edition. Crown 8vo. cloth,  
10s. 6d.
3. **A Treatise on the Differential Calculus :**  
With numerous Examples. Second Edition. Crown 8vo. cloth,  
10s. 6d.
4. **A Treatise on the Integral Calculus,**  
And its Applications; with numerous Examples. Crown 8vo.  
cloth, 10s. 6d.
5. **Examples of Analytical Geometry of Three**  
Dimensions. Crown 8vo. cloth, 4s.
6. **A Treatise on Analytical Statics.**  
With numerous Examples. Second Edition. Crown 8vo. cloth,  
10s. 6d. *Nearly Ready.*



**EXAMPLES OF**  
**ANALYTICAL GEOMETRY**  
**OF THREE DIMENSIONS.**



**Cambridge :**

**PRINTED BY C. J. CLAY, M.A.  
AT THE UNIVERSITY PRESS.**

EXAMPLES OF  
ANALYTICAL GEOMETRY  
OF THREE DIMENSIONS.

COLLECTED BY


I. TODHUNTER, M.A.

FELLOW AND ASSISTANT TUTOR OF ST JOHN'S COLLEGE, CAMBRIDGE.

Cambridge :  
MACMILLAN AND CO.  
1858.

[The right of Translation is reserved.]


183. G. 10.  
~~S. 1. 1. 10.~~



A COLLECTION of examples in illustration of Analytical Geometry of Three Dimensions has long been required both by students and teachers, and the present work is published with the view of supplying the want. These examples have been principally obtained from University and College Examination papers, but many of them are original. The results of the examples are given at the end of the book, together with hints for the solution of some of them.

ST JOHN'S COLLEGE,

July 17th, 1858.



## EXAMPLES OF ANALYTICAL GEOMETRY OF THREE DIMENSIONS.

### I. *The straight line and plane.*

Let  $OA$ ,  $OB$ ,  $OC$  be three edges of a rectangular parallelepiped which meet in a point  $O$ ; take  $O$  for the origin and the directions of  $OA$ ,  $OB$ ,  $OC$  for the axes of  $x$ ,  $y$ ,  $z$  respectively; complete the parallelepiped; let  $D$  be the vertex opposite to  $A$ ,  $E$  that opposite to  $B$ ,  $F$  that opposite to  $C$ ,  $G$  that opposite to  $O$ . Let  $OA = a$ ,  $OB = b$ ,  $OC = c$ ; and use these data in the following Examples from 1 to 11.

1. Find the equation to the plane passing through  $D$ ,  $E$ ,  $F$ .
2. Find the equation to the plane passing through  $G$ ,  $A$ ,  $B$ .
3. Find the equation to the plane passing through  $G$ ,  $O$ ,  $A$ ; also the equation to the plane passing through  $G$ ,  $O$ ,  $B$ .
4. Find the equations to the line  $OG$ .
5. Find the equations to the lines  $EB$  and  $AD$ .
6. Find the length of the perpendicular from the origin on the plane in Example 1.
7. Find the length of the perpendicular from  $C$  on the plane in Example 2.
8. Find the angle between the planes in Example 3.
9. Find the angle between the line in Example 4, and the normal to the plane in Example 1.
10. Find the angle between the lines in Example 5.

11. Find the equations to the line which passes through  $O$ , and the centre of the face  $AEGF$ .

12. Interpret the equation

$$x^2 + y^2 + z^2 = (x \cos \alpha + y \cos \beta + z \cos \gamma)^2,$$

$$\text{where } \cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1.$$

13. Find the angle between the line  $\frac{x}{2} = \frac{y}{\sqrt{3}} = \frac{z}{\sqrt{2}}$ , and the line  $\frac{x}{\sqrt{3}} = y, z = 0$ .

14. Find the equation to the plane which contains the line whose equations are

$$Ax + By + Cz = D, \quad A'x + B'y + C'z = D',$$

$$\text{and the point } (\alpha, \beta, \gamma).$$

15. Find the equation to the plane which passes through the origin and through the line of intersection of the planes

$$Ax + By + Cz = D, \quad \text{and } A'x + B'y + C'z = D';$$

and determine the condition that it may bisect the angle between them.

16. Find the equation to the plane which passes through the two parallel lines

$$\frac{x-a}{l} = \frac{y-b}{m} = \frac{z-c}{n}; \quad \frac{x-a'}{l} = \frac{y-b'}{m} = \frac{z-c'}{n}.$$

17. The equation to *one* plane through the origin bisecting the angle between the lines through the origin, the direction cosines of which are  $l_1, m_1, n_1$ , and  $l_2, m_2, n_2$ , and perpendicular to the plane containing them is

$$(l_1 - l_2)x + (m_1 - m_2)y + (n_1 - n_2)z = 0;$$

and the equation to the *other* plane is

$$(l_1 + l_2)x + (m_1 + m_2)y + (n_1 + n_2)z = 0.$$

18. Shew that the equation to a plane which passes through the point  $(\alpha, \beta, \gamma)$  and cuts off portions  $a, b$  from the axes of  $x$  and  $y$  respectively is

$$\frac{x}{a} + \frac{y}{b} - 1 = \frac{z}{\gamma} \left\{ \frac{\alpha}{a} + \frac{\beta}{b} - 1 \right\}.$$

19. Find the equation to the plane which contains a given line and is perpendicular to a given plane.

20. Shew that if the straight lines

$$\frac{x}{a} = \frac{y}{\beta} = \frac{z}{\gamma}, \quad \frac{x}{a^2\alpha} = \frac{y}{b^2\beta} = \frac{z}{c^2\gamma}, \quad \frac{x}{l} = \frac{y}{m} = \frac{z}{n},$$

lie in one plane, then

$$\frac{l}{a} (b^2 - c^2) + \frac{m}{\beta} (c^2 - a^2) + \frac{n}{\gamma} (a^2 - b^2) = 0.$$

21. Find the length of the perpendicular from the point  $(1, -1, 2)$  upon the line  $x = y = 2z$ .

22. Find the equation to the plane which passes through the lines

$$\frac{x-a}{l} = \frac{y-b}{m} = \frac{z-c}{n},$$

$$\text{and } \frac{x-a}{l'} = \frac{y-b}{m'} = \frac{z-c}{n'}.$$

23. Find the equations to a straight line which passes through the point  $(a, b, c)$  and makes a given angle with the plane

$$Ax + By + Cz = 0.$$

24. Find the equation to the plane perpendicular to a given plane, such that their line of intersection shall lie in one of the co-ordinate planes.

25. If the three adjacent edges of a cube be taken for the co-ordinate axes, find the co-ordinates of the points in which



a plane perpendicular to the diagonal through the origin and bisecting that diagonal will meet the edges.

26. Determine the plane which contains a given straight line, and makes a given angle with a given plane.

27. A straight line makes an angle of  $60^\circ$  with one axis, and an angle of  $45^\circ$  with another; what angle does it make with the third axis?

28. Interpret  $x^2 = y^2 = z^2$ .

29. Find the condition that must hold in order that the equations

$$x = cy + bz, \quad y = az + cx, \quad z = bx + ay$$

may represent a straight line; and shew that the equations to the line then are

$$\frac{x}{\sqrt{(1-a^2)}} = \frac{y}{\sqrt{(1-b^2)}} = \frac{z}{\sqrt{(1-c^2)}}.$$

30. Through the origin and the line of intersection of the planes

$$x \cos \alpha + y \cos \beta + z \cos \gamma - p = 0,$$

$$\text{and } x \cos \alpha_1 + y \cos \beta_1 + z \cos \gamma_1 - p_1 = 0,$$

a plane is drawn; perpendicular to this plane and through its line of intersection with the plane of  $(x, y)$  another plane is drawn; find its equation.

31. Find the equation to the plane which passes through a given point and is perpendicular to the line of intersection of two given planes.

32. From any point  $P$  are drawn  $PM, PN$  perpendicular to the planes of  $(z, x)$  and  $(z, y)$ ; if  $O$  be the origin,  $\alpha, \beta, \gamma, \theta$  the angles which  $OP$  makes with the co-ordinate planes and the plane  $OMN$ , then will

$$\operatorname{cosec}^2 \theta = \operatorname{cosec}^2 \alpha + \operatorname{cosec}^2 \beta + \operatorname{cosec}^2 \gamma.$$

33. Apply the equation to a plane

$$x \cos \alpha + y \cos \beta + z \cos \gamma = \delta$$

to prove the following theorem; a triangle is projected on each of three rectangular planes; shew that the sum of the pyramids which have these projections for bases and a common vertex in the plane of the triangle is equal to the pyramid which has the triangle for base and the origin for vertex.

34. Find the equations to the straight line joining the points  $(a, b, c)$  and  $(a', b', c')$ ; and shew that it will pass through the origin if  $aa' + bb' + cc' = \rho\rho'$ , where  $\rho$  and  $\rho'$  are the distances of the points respectively from the origin.

35. Express the equations to the line

$$a_1x + b_1y + c_1z = a_2x + b_2y + c_2z = a_3x + b_3y + c_3z$$

in the form

$$\frac{x}{l} = \frac{y}{m} = \frac{z}{n}.$$

36. Shew that the equations

$$\frac{x^3 + 1}{x + 1} = \frac{y^3 + 1}{y + 1} = \frac{z^3 + 1}{z + 1}$$

represent four straight lines, and that the angle between any two of them  $= \cos^{-1} \left( \frac{1}{3} \right)$ .

37. The equations to two planes are

$$lx + my + nz = p, \quad l'x + m'y + n'z = p',$$

where

$$l^2 + m^2 + n^2 = 1, \quad l'^2 + m'^2 + n'^2 = 1;$$

find the lengths of the perpendiculars from the origin upon the two planes which pass through their line of intersection and bisect the angles between them.

38. Find the equation to a plane parallel to two given straight lines; hence determine the shortest distance between two given straight lines.

39. Find the equation to a plane which passes through two given points and is perpendicular to a given plane.

40. There are  $n$  planes of which no two are parallel to each other, no three are parallel to the same right line, and no four pass through the same point; prove that the number of lines of intersection of the planes is  $\frac{n}{2}(n-1)$ , and that the number of points of intersection of those lines is

$$\frac{n(n-1)(n-2)}{1 \cdot 2 \cdot 3}.$$

41. Find the shortest distance between the point  $(\alpha, \beta, \gamma)$  and the plane

$$Ax + By + Cz = D.$$

42. Find the equation to the plane which passes through the origin and makes equal angles with three given straight lines which pass through the origin.

43. Determine the co-ordinates of the point which divides in a given ratio the distance between two points.

Hence shew that the equation

$$Ax + By + Cz = D$$

must represent a *plane*, according to Euclid's definition of a plane.

44. Three planes meet in a point, and through the line of intersection of each pair a plane is drawn perpendicular to the third; prove that in general the planes thus drawn pass through the same line.

45. The equations to a line are

$$\frac{\alpha - cy + bz}{a} = \frac{\beta - az + cx}{b} = \frac{\gamma - bx + ay}{c};$$

express them in the ordinary symmetrical form.

46. Find the condition that must subsist in order that the equations

$$a + mz - ny = 0, \quad b + nx - lz = 0, \quad c + ly - mx = 0$$

may represent a straight line; and supposing this condition to be satisfied put the equations in the ordinary symmetrical form.

47. Supposing the equations in the preceding question to represent a straight line, find in a symmetrical form the equations to the line from the origin perpendicular to the given line; also determine the co-ordinates of the point of intersection.

48. The locus of the middle points of all straight lines parallel to a fixed plane and terminated by two fixed right lines which do not intersect is a straight line.

49. The equation to a plane is  $lx + my + nz = 0$ ; find the equations to a line lying in this plane and bisecting the angle formed by the intersections of the given plane with the co-ordinate planes of  $(z, x)$  and  $(z, y)$ .

50. A straight line, whose equations are given, intersects the co-ordinate planes in three points; find the angles included between the lines which join these points with the origin; and if these angles  $(\alpha, \beta, \gamma)$  be given, shew that the equation to the surface traced out by the line in all positions is

$$x \sqrt{\tan \alpha} + y \sqrt{\tan \beta} + z \sqrt{\tan \gamma} = 0.$$

II. *Surfaces of the second order.*

51. If the normal  $n$  at any point of an ellipsoid terminated in the plane of  $(x, y)$  make angles  $\alpha, \beta, \gamma$  with the semi-axes  $a, b, c$ , and  $p$  be the perpendicular from the centre on the tangent plane, then

$$n \cdot p = c^2, \quad \text{and} \quad p^2 = a^2 \cos^2 \alpha + b^2 \cos^2 \beta + c^2 \cos^2 \gamma.$$

52. Find a point on an ellipsoid such that the tangent plane cuts off equal intercepts from the axes. Also find a point such that the intercepts are proportional to the axes.

53. If  $a, b, c$  be the semi-axes of an ellipsoid taken in order, and  $e, e'$  the excentricities of the principal sections containing the mean axis, shew that the perpendiculars from the centre on the tangent planes at every point of the section of the surface made by the plane  $abe'z = c^2ex$  are equal.

54. From a given point  $O$ , a line  $OP$  is drawn meeting a given plane in  $Q$ , and the rectangle  $OP \cdot OQ$  is invariable; find the locus of  $P$ .

55. Sections of an ellipsoid are made by planes which all contain the least axis; find the locus of the foci of the sections.

56. Find the locus of a point which is equidistant from every point of the circle determined by the equations

$$x^2 + y^2 + z^2 = a^2, \quad lx + my + nz = p.$$

57. Shew that the section of the surface  $z^2 = xy$  by the plane  $z = x + y + c$  is a circle.

58. Interpret the equation

$$x^2 + y^2 + z^2 = (lx + my + nz)^2.$$

59.  $A, B, C$  are three fixed points, and  $P$  a point in space such that  $PA^2 + PB^2 = PC^2$ ; find the locus of  $P$ , and explain the result when  $ACB$  is a right or obtuse angle.

60. Shew that the equation  $\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$ , where  $a, b, c$  are in order of magnitude, may be written thus,

$$k^2(x^2 + y^2 + z^2 - b^2) - x^2 + m^2 z^2 = 0,$$

where  $k$  and  $m$  are certain constants. Hence shew that two circular sections of an ellipsoid can be obtained by cutting it by planes passing through its mean axis.

61. A sphere ( $C$ ) and a plane are given; shew that if any sphere ( $C'$ ) be described touching the plane at a given point and cutting  $C$ , the plane of section always contains a given right line. Shew also that if the point of contact be not given, and if the plane of section always contain a given point, the centre of the sphere  $C'$  will always be upon a given paraboloid.

62. If  $A, B, C$  be extremities of the axes of an ellipsoid, and  $AC, BC$  be the principal sections containing the least axis, find the equations to the two cones whose vertices are  $A, B$ , and bases  $BC, AC$  respectively; shew that the cones have a common tangent plane, and a common parabolic section, the plane of the parabola and the tangent plane intersecting the ellipsoid in ellipses, the area of one of which is double that of the other; and if  $l$  be the latus rectum of the parabola,  $l_1, l_2$  of the sections  $AC, BC$ , prove that

$$\frac{1}{l^2} = \frac{1}{l_1^2} + \frac{1}{l_2^2}.$$

63. Tangent planes are drawn to an ellipsoid from a given external point; find the equation to the cone which has its vertex at the origin and passes through all the points of contact of the tangent planes with the ellipsoid.

64. If tangent planes be drawn to an ellipsoid from any point in a plane parallel to that of  $(x, y)$ , the curve which contains all the points of contact will lie in a plane which always cuts the axis of  $z$  in the same point.

65. Shew that the tangent plane to an ellipsoid is expressed by the equation

$$lx + my + nz = \sqrt{(l^2 a^2 + m^2 b^2 + n^2 c^2)}.$$

66. Form the equation to the plane which passes through a given point of an ellipsoid, through the normal at that point, and through the centre of the ellipsoid.

67. A line passing through a given point moves so that the projection of any portion of it on a given line bears a constant ratio to the length of that portion; find the equation to the surface which it traces out.

68. Find the length of the perpendicular from a given point on a given straight line in space.

Investigate the equation to a right cone, having the axis, vertex, and vertical angle given; and determine the condition under which the section made by a plane parallel to one of the co-ordinate planes will be an ellipse.

69. Determine the radii of the spheres which touch the co-ordinate planes and the plane  $x + y + z = h$ .

70. An ellipsoid is intersected in the same curve by a variable sphere, and a variable cylinder; the cylinder is always parallel to the least axis of the ellipsoid, and the centre of the sphere is always at the focus of a principal section containing this axis. Prove that the axis of the cylinder is invariable in position, and that the area of its transverse section varies as the surface of the sphere.

71. Three edges of a tetrahedron, in length  $a$ ,  $b$ , and  $c$ , are mutually at right angles; prove that if these three edges be taken as axes, the equation to the cone which has the origin for vertex, and for its base the circle circumscribed about the opposite face, is

$$\left(\frac{b}{c} + \frac{c}{b}\right)yz + \left(\frac{c}{a} + \frac{a}{c}\right)zx + \left(\frac{a}{b} + \frac{b}{a}\right)xy = 0,$$

and that the plane  $ax + by + cz = 0$  is parallel to the sub-conary sections of the cone.

Find the corresponding equations when either of the other angular points of the tetrahedron is taken as vertex.

72. Tangent planes to the surface whose equation is

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$$

pass through a point  $P$ ; prove that a sphere can be described through the curve of contact provided  $P$  be on a certain right line passing through the origin.

73. A sphere touches each of two right lines which are inclined to each other at a right angle but do not meet; shew that the locus of its centre is an hyperbolic paraboloid.

74. Shew that by properly choosing the rectangular axes any two lines may be represented by the equations

$$\left. \begin{array}{l} y = mx \\ z = c \end{array} \right\}, \quad \left. \begin{array}{l} y = -mx \\ z = -c \end{array} \right\}.$$

Determine the locus of a point which moves so as always to be equally distant from two given straight lines.

75. A tangent plane to an ellipsoid includes between itself and the co-ordinate planes a constant volume; find the locus of the points of contact.

76. A tangent plane to an ellipsoid passes through a fixed point in one of the axes produced; find the point of contact when the volume between this plane and the co-ordinate planes is a minimum.

77. Prove that the locus of a point whose distance from a fixed point is always in a given ratio to its distance from a fixed line is a surface of revolution of the second degree.

78. Two systems of rectangular axes have the same origin; if a plane, or an ellipsoid whose centre is the origin, cut them at distances  $a, b, c$ , or  $a', b', c'$ , respectively, then

$$\frac{1}{a^2} + \frac{1}{b^2} + \frac{1}{c^2} = \frac{1}{a'^2} + \frac{1}{b'^2} + \frac{1}{c'^2}.$$



79. If a moveable line make with any number of fixed lines the angles  $\theta_1, \theta_2, \theta_3, \dots$  so that

$$a_1 \cos \theta_1 + a_2 \cos \theta_2 + a_3 \cos \theta_3 + \dots = \text{a constant},$$

the lines all passing through a fixed point; shew that the moveable line will always lie on the surface of a right cone.

80. Two prolate spheroids have a common focus. If their surfaces meet, the points of intersection will lie in planes which pass through the line common to the directrix planes.

81. A cone passes through the principal section  $(b, c)$  of an ellipsoid; shew that the other section is a curve in a plane perpendicular to that of  $(b, c)$  if the vertex of the cone be any point of the surface

$$\frac{z^2}{c^2} + \frac{y^2}{b^2} - \frac{x^2}{a^2} = 1.$$

82. An ellipsoid is cut by a plane parallel to one of its principal planes; shew that all normals to the ellipsoid at points in the curve of section pass through two straight lines situated in the other principal planes.

83. An ellipsoid is constantly touched in two points by a sphere of given radius whose centre moves in one of its principal planes; find the locus of the centre of the sphere, and shew that it will describe a portion of an equilateral hyperbola, if the plane in which it moves be that which contains the greatest and the least axes, and if the foci of the other principal sections be equidistant from the centre of the ellipsoid.

84. An elliptic cylinder is cut by a plane passing through the straight line which touches the base at the extremity of its axis minor; shew that, if the sine of the inclination of the cutting plane to the base do not exceed the excentricity of the base, the locus of the foci of the section will be a circle.

85. Find the curves of intersection of the surface

$$xy + yz + zx = a^2$$

with planes parallel to  $x + y + z = 0$ .

86. If a sphere be placed in a paraboloid of revolution, shew that the section of the paraboloid made by any plane drawn touching the sphere is a conic section having the point of contact for a focus.

87. Shew that any plane parallel to the planes of  $(x, z)$  or  $(y, z)$  will meet the surface whose equation is  $xy = az$  in a straight line; if a plane be made to pass through that line and touch the surface will the plane touch the surface in every point of the line?

88. Through any point of the curve of intersection of the surfaces

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} - \frac{z^2}{c^2} = 1, \quad x^2 + y^2 + z^2 = a^2 + b^2 - c^2,$$

two straight lines can be drawn on the former surface at right angles to one another.

89. Prove that the plane  $z = mx + ny$  will cut the surface

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = \frac{z^2}{c^2}$$

in two straight lines which are at right angles to one another if

$$\frac{1}{b^2} + \frac{1}{a^2} + m^2 \left( \frac{1}{b^2} - \frac{1}{c^2} \right) + n^2 \left( \frac{1}{a^2} - \frac{1}{c^2} \right) = 0.$$

90. Shew that an infinite number of straight lines may be drawn lying on the surface whose equation is

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = \frac{4z}{c}.$$

If two such lines be drawn through a point in the plane of  $(x, z)$  whose co-ordinates are  $(x', z')$  the angle between them is

$$\cos^{-1} \frac{a^2 - b^2 + cz'}{a^2 + b^2 + cz'}.$$

91. Prove that the points on the surface  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = \frac{4z}{c}$ , the straight lines through which coincident with the surface are at right angles to each other, lie in a plane parallel to the plane of  $(x, y)$  and at a distance from it equal to  $\frac{b^2 - a^2}{c}$ .

92. Determine the condition necessary in order that the plane  $lx + my + nz = 0$  may cut the cone  $ayz + bzx + cxy = 0$  in two straight lines at right angles to each other; and shew that in that case the straight line through the origin perpendicular to the plane will also lie on the cone.

93. Find the relations between the coefficients of the equation

$$ax^2 + by^2 + cz^2 + 2a'yz + 2b'xz + 2c'xy + 2a''x + 2b''y + 2c''z = f,$$

that it may represent a surface of revolution.

$$\text{If } \frac{b'c'}{a'} - a = 0, \quad \frac{c'a'}{b'} - b = 0, \quad \frac{a'b'}{c'} - c = 0,$$

what is the nature of the surface?

94. Determine the conditions that the equation

$$ax^2 + by^2 + cz^2 + 2a'yz + 2b'zx + 2c'xy = 0$$

may represent a right circular cone; obtaining them in the form

$$\frac{a'b'}{c'} + \frac{a^2 - b^2}{a - b} = 0, \quad \frac{c'a'}{b'} + \frac{c^2 - a^2}{c - a} = 0.$$

95. Find the relations between the coefficients of the equation

$$ax^2 + by^2 + cz^2 + 2b'xz + 2c'xy + 2a''x + 2b''y + 2c''z = f,$$

that it may represent a surface of revolution.

96. Prove that the right circular cylinders described about the ellipsoid

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1,$$

$b$  being the mean axis, are represented by the equation

$$(b^2 - c^2)x^2 - (c^2 - a^2)y^2 + (a^2 - b^2)z^2$$

$$\pm 2\sqrt{(a^2 - b^2)}\sqrt{(b^2 - c^2)}xz = (a^2 - c^2)b^2.$$

97. Find the locus of the vertices of all right cones which have the same given ellipse as base.

98. The semiaxes of an ellipsoid are  $a, b, c$ ; if planes are drawn through the centre to cut it in sections which have the constant area  $\frac{\pi abc}{f}$ , shew that the normals to these planes will all lie upon the cone

$$x^2 (a^2 - f^2) + y^2 (b^2 - f^2) + z^2 (c^2 - f^2) = 0.$$

99. On every central radius vector of the ellipsoid

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1,$$

whose centre is  $C$  is taken a point  $P$  such that  $\pi k CP$  is the area of the section of the ellipsoid made by the plane through  $C$  perpendicular to  $CP$ , where  $k$  is a line of constant length; shew that the equation to the locus of  $P$  is

$$a^2 x^2 + b^2 y^2 + c^2 z^2 = \frac{a^2 b^2 c^2}{k^2}.$$

100. The sum of the squares of the reciprocals of the areas of the sections of an ellipsoid made by any three diametral planes at right angles to each other is constant.

101. An ellipsoid and hyperboloid are concentric and confocal; if a tangent plane be drawn to the asymptotic cone of the hyperboloid, the section of the ellipsoid will be of constant area.

102. A sphere and an ellipsoid which intersect are described about the same point as centre; prove that the product of the areas of the greatest and least sections of the ellipsoid, made by planes passing through the centre and any point of the line of intersection of the two surfaces will be constant.

103. Find the area of the section of the ellipsoid

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1,$$

made by the plane  $lx + my + nz = \delta$ .

104. Shew that the sections of a surface of the second order made by parallel planes are similar curves. Having given the area of the section of an ellipsoid by the plane  $lx + my + nz = 0$ , find the area of the section made by the plane  $lx + my + nz = \delta$ .

105. If  $S$  be the area of a section of an ellipsoid made by a plane at the distance  $h$  from the centre,  $S'$  that of the parallel section through the centre, and  $p$  the perpendicular from the centre on the parallel tangent plane, shew that

$$S = S' \left( 1 - \frac{h^2}{p^2} \right).$$

106. Tangent planes are drawn to an ellipsoid from a given point; shew that an ellipsoid similar to the given ellipsoid and similarly situated can be made to pass through the given point, the points of contact, and the centre of the given ellipsoid.

107. Normals are drawn to the ellipsoid

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$$

at the points where it is intersected by the plane  $z = h$ . Shew that the locus of the intersection of these normals with the plane of  $(x, y)$  is the ellipse

$$\frac{x^2 a^2}{(a^2 - c^2)^2} + \frac{y^2 b^2}{(b^2 - c^2)^2} = 1 - \frac{h^2}{c^2}.$$

108. All the normals to an ellipsoid meet the plane of  $(x, y)$  within an ellipse whose equation is

$$\frac{a^2 x^2}{(a^2 - c^2)^2} + \frac{b^2 y^2}{(b^2 - c^2)^2} = 1.$$

Determine what is represented by the following equations.

109.  $5y^2 - 2x^2 - z^2 + 4xy - 6yz + 8xz - 1 = 0.$

110.  $2y^2 - 5x^2 + 2z^2 + 10xy + 4yz + 4y + 16z + 18 = 0,$

111.  $4y^2 - 9x^2 + 2xy + 36x - 8y - 4z - 32 = 0.$

112.  $y^2 - 4xy + 4x^2 - 6x + 3z = 0.$

113.  $x^2 - y^2 + z^2 - 4xy + 6xz - 2yz = f.$

114.  $(ay - bx)^2 + (cx - az)^2 + (bz - cy)^2 = f.$

115.  $xy + xz + yz - 2x + 6y - 8z = f.$

116.  $xy + yz + xz = a^2.$

117. Find the centre of the surface

$$(ay + \beta x - cy)^2 = a\beta (xy - z^2),$$

118. Find the centre of the surface

$$x^2 + y^2 + z^2 + 4xy - 2xz - 4yz + 2x + 4y - 2z = 0.$$

119. Find the centre of the surface

$$x^2 + 2y^2 + 3z^2 + 2(xy + yz + xz) + x + y + z = 1.$$

120. Shew that the equation  $yz + zx + xy = a^2$  may be reduced to

$$2x^2 - (y^2 + z^2) = 2a^2$$

by transforming the axes.

121. Shew that the equation

$$x^2 + y^2 + z^2 + xy + yz + zx = a^2$$

represents an oblate spheroid whose polar axis is to its equatorial in the ratio of 1 to 2, and the equations to whose axis are  $x = y = z,$

122. What is represented by the following equation?

$$x^2 + y^2 + z^2 + k(xy + yz + zx) = f.$$

123. If two concentric surfaces of the second order have the same foci for their principal sections they will cut one another everywhere at right angles.

124. Find the locus of the intersection of three planes at right angles to each other, each of which touches one of the following three ellipsoids,

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1, \quad \frac{x^2}{a^2 + h^2} + \frac{y^2}{b^2 + h^2} + \frac{z^2}{c^2 + h^2} = 1,$$

$$\frac{x^2}{a^2 + k^2} + \frac{y^2}{b^2 + k^2} + \frac{z^2}{c^2 + k^2} = 1.$$

125. Determine the position of the circular sections of an hyperboloid of two sheets, and shew that the same plane will cut the asymptotic cone in a circle.

126. If  $x_1, y_1, z_1; x_2, y_2, z_2; x_3, y_3, z_3$  be the co-ordinates of the extremities of a set of conjugate diameters of an ellipsoid, shew that

$$x_1^2 + x_2^2 + x_3^2 = a^2, \quad y_1^2 + y_2^2 + y_3^2 = b^2, \quad z_1^2 + z_2^2 + z_3^2 = c^2,$$

$$x_1 y_1 + x_2 y_2 + x_3 y_3 = 0, \quad x_1 z_1 + x_2 z_2 + x_3 z_3 = 0, \quad y_1 z_1 + y_2 z_2 + y_3 z_3 = 0.$$

127. If spheres be described on three semi-conjugate diameters of an ellipsoid as diameters, the locus of their intersection is the surface determined by

$$a^2 x^2 + b^2 y^2 + c^2 z^2 = 3(x^2 + y^2 + z^2)^2.$$

128. A plane is drawn through the extremities of three semi-conjugate diameters of an ellipsoid; find the locus of the intersection of this plane with the perpendicular on it from the centre.

129. Tangent planes at the extremities of three conjugate diameters of an ellipsoid intersect in the ellipsoid whose equation is

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 3.$$

130. A prolate spheroid is cut by any plane through one of its foci; prove that the focus is a focus of the section.

131. Shew that the locus of the diameters of the ellipsoid

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1,$$

which are parallel to the chords bisected by tangent planes to the cone

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} - \frac{z^2}{c^2} = 0, \text{ is the cone } \frac{a^2 x^2}{a^4} + \frac{b^2 y^2}{b^4} - \frac{\gamma^2 z^2}{c^4} = 0,$$

132. If three straight lines at right angles to each other touch the ellipsoid

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1,$$

and intersect each other in the point  $(x', y', z')$  shew that

$$x'^2 (b^2 + c^2) + y'^2 (c^2 + a^2) + z'^2 (a^2 + b^2) = b^2 c^2 + c^2 a^2 + a^2 b^2.$$

133. Find the greatest angle between the normal at any point of an ellipsoid, and the central radius vector at that point.

134. If four similar and similarly situated surfaces of the second order intersect each other, the planes of their intersections two and two all pass through one point.

135. If three chords be drawn mutually at right angles through a fixed point within a surface of the second order whose equation is  $u = 0$ , shew that  $\Sigma \frac{1}{R \cdot r}$  will be constant where  $R$  and  $r$  are the two portions into which any one of the chords drawn through the fixed point is divided by that point.

Prove also that the same will be true if instead of the fixed point there be substituted any point on the surface whose equation is  $u = c$ .

136. Let  $\frac{x}{l} = \frac{y}{m} = \frac{z}{n}$  be the equations to a straight line; find the equation to a surface every point of which is at the same distance from this line as from the point  $(a, \beta, \gamma)$ ; and shew that the plane  $lx + my + nz = \delta$  cuts the surface in a straight line.



137.  $A$  and  $B$  are two similar and concentric ellipsoids, the homologous axes being in the same straight line;  $C$  is a third ellipsoid similar to either of the former, its centre being on the surface of  $B$ , and axes parallel to those of  $A$  or  $B$ ; shew that the plane of intersection of  $A$  and  $C$  is parallel to the tangent plane to  $B$  at the centre of  $C$ .

138. If a parallelopiped be inscribed in an ellipsoid its edges will be parallel to a system of conjugate diameters.

139. The edges of a parallelopiped are  $2a, 2b, 2c$ ; shew that an ellipsoid concentric with it and whose semidiameters parallel to the edges are  $a\sqrt{2}, b\sqrt{2}, c\sqrt{2}$ , intersects the faces in ellipses which touch each other and the edges.

140. Two similar and similarly situated ellipsoids are cut by a series of ellipsoids similar and similarly situated to the two given ones, so that the planes of intersection of any one of the series with each of the given ellipsoids make a right angle with one another. Shew that the centres of the series of ellipsoids lie on another ellipsoid,

141. If pyramids be formed between three conjugate diametral planes of an ellipsoid and a tangent plane, so that the products of the intercepted portions of the three conjugate diameters may be the least possible, the volumes of all these pyramids will be equal.

142. If from any point in an ellipsoid three straight lines are drawn mutually at right angles, prove that the plane which passes through their intersections with the surface passes through a point which is fixed so long as the original point is fixed. And shew that if the position of the original point on the surface is changed the locus of the point is an ellipsoid whose semiaxes are

$$\frac{a^2 - 2k^2}{a}, \frac{b^2 - 2k^2}{b}, \frac{c^2 - 2k^2}{c},$$

where

$$\frac{1}{k^2} = \frac{1}{a^2} + \frac{1}{b^2} + \frac{1}{c^2}.$$

143. If two right circular cones whose axes are parallel intersect, the projection of their curve of intersection on the plane passing through the axes is a parabola.

144. An ellipsoid is cut by a plane the distance of which from the centre bears a constant ratio to its distance from the parallel tangent plane; shew that the volume of the cone whose base is the section and vertex the centre of the ellipsoid is invariable.

145. Shew that by properly choosing oblique axes the equation to an ellipsoid may be put in the form

$$x^2 + y^2 + z^2 = a'^2;$$

shew that an infinite number of systems of suitable axes can be found.

146. Determine whether the equation to an hyperboloid of one sheet can always be put in the form

$$x^2 + y^2 - z^2 = a'^2,$$

by properly choosing oblique axes.

147. A straight line revolving round an axis which it does not meet generates an hyperboloid.

148. Prove that the equations to the principal axes of the surface

$$ax^2 + by^2 + cz^2 + 2a'yz + 2b'xz + 2c'xy = 1$$

are

$$x \{a'(\mu + a) - b'c'\} = y \{b'(\mu + b) - c'a'\} = z \{c'(\mu + c) - a'b'\};$$

$\mu$  being determined by the equation

$$\begin{aligned} &(\mu + a)(\mu + b)(\mu + c) - a'^2(\mu + a) - b'^2(\mu + b) \\ &\quad - c'^2(\mu + c) + 2a'b'c' = 0. \end{aligned}$$

149. In an ellipsoid the sum of the lengths of three conjugate diameters is greatest when the diameters are equal.

150. A surface of the second order circumscribes a tetrahedron and each face is parallel to the tangent plane at the opposite angular point; shew that the equation to the surface, taking as oblique axes the three edges of the tetrahedron which meet in one of its angular points, is

$$\frac{x^2 - ax}{a^2} + \frac{y^2 - by}{b^2} + \frac{z^2 - cz}{c^2} + \frac{1}{abc} \{ayz + bzx + cxy\} = 0.$$

151. Shew that the centre of the surface in the preceding question coincides with the centre of gravity of the tetrahedron.

152. If two hyperboloids of one sheet have their corresponding axes equal and parallel, shew that they intersect in a plane curve, the plane of which is parallel to the diametral planes conjugate to the line joining their centres; and find where it cuts this line.

153. Shew that any plane which contains two parallel generating lines of an hyperboloid of one sheet passes through the centre of the surface.

154. The points on an ellipsoid, the normals at which intersect the normal at a fixed point, lie on a cone of the second order whose vertex is the fixed point.

155. Normals are drawn to an hyperboloid of one sheet at points lying along a generating line; shew that these normals are all parallel to a fixed plane.

156. If one of the co-ordinates of an ellipsoid be produced so that the part produced equals the sum of the other two, its extremity will trace out a concentric ellipsoid whose volume is equal to that of the original ellipsoid.

157. Investigate the conditions necessary in order that the equation

$$ax^2 + by^2 + cz^2 + 2a'yz + 2b'zx + 2c'xy = 0$$

may represent two planes.

158. The equation to a surface of the second order is

$$ax^2 + by^2 + cz^2 + 2a'yz + 2b'zx + 2c'xy \\ + 2a''x + 2b''y + 2c''z + f = 0;$$

shew that the diametral planes will be parallel to a given line if

$$aa'^2 + bb'^2 + cc'^2 - abc - 2a'b'c' = 0,$$

and to a given plane if

$$a^2 - bc = 0, \quad b^2 - ac = 0, \quad c^2 - ab = 0.$$

159. Find the line in which the two surfaces intersect whose equations are

$$x^2 \cos 2\theta_1 + y^2 \cos 2\theta_2 + z^2 \cos 2\theta_3 + 2xy \cos (\theta_1 + \theta_2) \\ + 2zx \cos (\theta_1 + \theta_3) + 2zy \cos (\theta_2 + \theta_3) = 0, \\ x^2 \sin 2\theta_1 + y^2 \sin 2\theta_2 + z^2 \sin 2\theta_3 + 2xy \sin (\theta_1 + \theta_2) \\ + 2zx \sin (\theta_1 + \theta_3) + 2zy \sin (\theta_2 + \theta_3) = 0.$$

160. Shew that the cones determined by

$$\frac{x(yb + zc - xa)}{A} = \frac{y(zc + xa - yb)}{B},$$

and 
$$\frac{y(zc + xa - yb)}{B} = \frac{z(xa + yb - zc)}{C},$$

intersect in the line determined by

$$\frac{x}{A(Bb + Cc - Aa)} = \frac{y}{B(Cc + Aa - Bb)} = \frac{z}{C(Aa + Bb - Cc)};$$

and find the other lines in which they intersect.

161. Find the excentricity of any section of a paraboloid of revolution in terms of the angle of inclination of the cutting plane to the axis.

162. If through any fixed point chords be drawn to an ellipsoid, the intersection of pairs of tangent planes at their extremities all lie in one plane.

163. Let  $S$  be a fixed point, and  $EF$ ,  $LM$  two fixed straight lines; take a point  $P$  such that if  $PQ$  be drawn perpendicular to  $EF$  to meet  $LM$  in  $Q$ , then  $PQ$  may bear an invariable ratio to  $SP$ ; shew that the locus of  $P$  will be a surface of the second order. Shew also that when  $SP$ ,  $PQ$  are unequal, a section of the surface by a plane perpendicular to  $EF$  is a circle, and that when  $SP$ ,  $PQ$  are equal the surface has no centre.

164. The equation  $\frac{x^2 + y^2}{a^2} - \frac{z^2}{c^2} = 1$ , by varying  $c$ , represents a series of spheroidal surfaces. If normals be drawn to the spheroids from a fixed point in the axis of  $z$ , their intersections with the surfaces will lie on the surface of a given sphere.

165. An ellipsoid and hyperboloid of two sheets have the same principal diameters, both in magnitude and position. Let any tangent plane be drawn to the ellipsoid at a point  $P$ ; this will intersect in general the hyperboloid in a plane curve, through which if there be drawn any three tangent planes to the latter surface they will meet in a point  $P'$  which lies on the ellipsoid. Also if  $P'$  be the point of contact of a second tangent plane to the ellipsoid,  $P$  will be the intersection of a second set of tangent planes to the hyperboloid, drawn as before.

166. Two spheres being given in magnitude and position, every sphere which intersects them in given angles will touch two other fixed spheres and cut another at right angles.

167. The surface of an ellipsoid is represented by the equations

$$x = a \cos \alpha, \quad y = b \cos \beta, \quad z = c \cos \gamma;$$

shew that the three lines whose direction angles are  $(\alpha, \beta, \gamma, \&c.)$  corresponding to any system of conjugate diameters are perpendicular to each other.

If  $CP_1$ ,  $CP_2$ ,  $CP_3$  be drawn perpendicular to three conjugate tangent planes

$$\frac{1}{CP_1^2} + \frac{1}{CP_2^2} + \frac{1}{CP_3^2} = \frac{1}{a^2} + \frac{1}{b^2} + \frac{1}{c^2};$$

and if they cut the ellipsoid in points  $Q_1, Q_2, Q_3$ , respectively,

$$\frac{1}{CP_1^2 CQ_1^2} + \frac{1}{CP_2^2 CQ_2^2} + \frac{1}{CP_3^2 CQ_3^2} = \frac{1}{a^4} + \frac{1}{b^4} + \frac{1}{c^4}.$$

168. A plane is drawn according to a certain assigned law cutting an ellipsoid; find the locus of the vertex of the cone which touches the ellipsoid in the curve of intersection.

If  $a, b, c$  be the semiaxes of the ellipsoid  $\frac{x}{a} + \frac{y}{\beta} + \frac{z}{\gamma} = 1$ , the equation to the cutting plane,  $\alpha, \beta, \gamma$ , being connected by the relation  $\frac{\alpha^4}{a^4} + \frac{\beta^4}{b^4} + \frac{\gamma^4}{c^4} = \text{constant}$ , the locus is a sphere.

169. Shew that the tangent lines to an ellipsoid from an external point touch it in a plane curve, and find the equations to the curve.

If the external point lies on a fixed plane, shew that the plane of contact passes through a fixed point; determine its co-ordinates.

If the external point lies on a fixed line, the plane of contact contains a fixed line; determine its equation.

170. If lines be drawn from any point of the surface  $\frac{x^2}{a^4} + \frac{y^2}{b^4} + \frac{z^2}{c^4} - \frac{1}{k^2} = 0$ , to touch the surface  $\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} - 1 = 0$ ; shew that the planes in which the curves of contact lie all touch the sphere  $x^2 + y^2 + z^2 = k^2$ .

171. If lines be drawn from the centre of an ellipsoid (whose semiaxes are  $a, b, c$ ) parallel to the generating lines of an enveloping cone, the conical surface  $S$  so formed will intersect the ellipsoid in two planes parallel to the plane of contact. The locus of the vertex of the enveloping cone which causes one of the planes to coincide with the plane of contact is

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 2;$$

and the locus of the vertices of all cones formed in the same manner as  $S$  which intersect the ellipsoid in plane curves, is

the straight line joining the centre of the ellipsoid and the vertex of the enveloping cone.

172. Three cylinders circumscribe a given ellipsoid; the axes of the cylinders are mutually at right angles; shew that the sum of the squares of the areas of sections of the cylinders by planes respectively perpendicular to their axes is constant.

173. Of all cones which envelope an ellipsoid, have their bases in the tangent plane at a given point  $P$ , and are of the same altitude, that is the least which has its vertex in the diameter through  $P$ ; and of all which have their vertices in this diameter, that is the least whose axis is twice that diameter.

174. If a globe be placed upon a table the breadth of the elliptic shadow cast by a candle (considered as a luminous point) will be independent of the position of the globe.

175. In the preceding example, if an ellipsoid having its least axis vertical be substituted for the globe, determine the condition of the shadow of the globe being circular. It may be shewn that the locus of the luminous point must be an hyperbola, and that the radius of the circular shadow is independent of the mean axis of the ellipsoid.

176. Of a series of cones enveloping an ellipsoid, the vertices lie on a concentric ellipsoid, similar to the given one and similarly situated. Prove that any two cones of the series intersect one another in two planes.

177. Prove that if, in the preceding example, the vertices are supposed to lie also on a third ellipsoid concentric with the other two and similarly situated, and whose axes are respectively as the squares of theirs, these two planes are at right angles to one another.

178. A cylinder is circumscribed about an ellipsoid, and at the extremities of the diameter parallel to the generating lines of the cylinder tangent planes are drawn: shew that the volume of all cylinders so shut in is constant.

179. The locus of the vertex of a cone which envelopes a given ellipsoid ( $A$ ) is a straight line passing through the

centre of ( $A$ ); an ellipsoid similar to  $A$  and similarly placed has the vertex of the cone for centre and cuts the cone in a curve ( $B$ ). If the major axis of this ellipsoid vary as the distance of its centre from  $A$ 's, prove that the locus of  $B$  is an elliptic cylinder.

180. Suppose a cylinder to envelope an ellipsoid, and suppose a tangent plane to be drawn to the ellipsoid at one extremity of the diameter which is parallel to the axis of the cylinder. Let a line be drawn from the centre of the ellipsoid to meet the ellipsoid, the above tangent plane, and the enveloping cylinder; and suppose  $r, s, t$  to denote the respective distances of the points of intersection from the centre of the ellipsoid. Shew that

$$\frac{1}{r^2} = \frac{1}{s^2} + \frac{1}{t^2}.$$

181. Suppose a cone to envelope an ellipsoid; let  $R'$  be the distance of the vertex from the centre of the ellipsoid,  $2R$  the length of the diameter of the ellipsoid which is in the direction of the line joining the centre of the ellipsoid with the vertex of the cone; and suppose a tangent plane to be drawn to the ellipsoid at one extremity of this diameter. Let a line be drawn from the centre of the ellipsoid to meet the ellipsoid, the above tangent plane, and the enveloping cone; and suppose  $r, s, t$  to denote the respective distances of the points of intersection from the centre of the ellipsoid. Shew that

$$\left(\frac{1}{R't} - \frac{1}{R's}\right)^2 = \left(\frac{1}{R'^2} - \frac{1}{R'^2}\right) \left(\frac{1}{r^2} - \frac{1}{s^2}\right).$$

182. Let  $\phi(x, y, z) = 0$  be the equation to a surface of the second order; if tangent lines be drawn to it from the point  $(\alpha, \beta, \gamma)$ , shew that the equation to the plane which contains all the points of contact is

$$2u + (x - \alpha) \frac{du}{d\alpha} + (y - \beta) \frac{du}{d\beta} + (z - \gamma) \frac{du}{d\gamma} = 0,$$

where  $u = \phi(\alpha, \beta, \gamma)$ .

183. Let  $\phi(x, y, z) = 0$  be the equation to a surface of



the second order; then the equation to the enveloping cone which has its vertex at the point  $(\alpha, \beta, \gamma)$  is

$$2u \left\{ (x-\alpha)^2 \frac{d^2u}{d\alpha^2} + (y-\beta)^2 \frac{d^2u}{d\beta^2} + (z-\gamma)^2 \frac{d^2u}{d\gamma^2} \right. \\ \left. + 2(x-\alpha)(y-\beta) \frac{d^2u}{d\alpha d\beta} + 2(y-\beta)(z-\gamma) \frac{d^2u}{d\beta d\gamma} + 2(z-\gamma)(x-\alpha) \frac{d^2u}{d\gamma d\alpha} \right\} \\ = \left\{ (x-\alpha) \frac{du}{d\alpha} + (y-\beta) \frac{du}{d\beta} + (z-\gamma) \frac{du}{d\gamma} \right\}^2,$$

where  $u = \phi(\alpha, \beta, \gamma)$ .

Ex. Determine the enveloping cone of the surface

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1, \text{ from the point } (a, b, c) \text{ as vertex.}$$

184. Let  $\phi(x, y, z) = 0$  be the equation to a surface of the second order; then the equation to the enveloping cylinder which has its generating lines parallel to the line  $\frac{x}{l} = \frac{y}{m} = \frac{z}{n}$  is

$$2u \left\{ l^2 \frac{d^2u}{dx^2} + m^2 \frac{d^2u}{dy^2} + n^2 \frac{d^2u}{dz^2} + 2lm \frac{d^2u}{dxdy} + 2mn \frac{d^2u}{dydz} + 2nl \frac{d^2u}{dzdx} \right\} \\ = \left( l \frac{du}{dx} + m \frac{du}{dy} + n \frac{du}{dz} \right)^2,$$

where  $u = \phi(x, y, z)$ .

Ex. Determine the enveloping cylinder of the surface  $\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$ , which has its generating lines parallel to the line  $\frac{x}{a} = \frac{y}{b} = \frac{z}{c}$ .

185. Shew that the surface

$$(ax + by + cz - 1)^2 + 2a'yz + 2b'zx + 2c'xy = 0$$

touches the co-ordinate axes; and find the equation to the

cone which has its vertex at the origin and passes through the curve of section of the surface by the plane through the points of contact.

186. Lines are drawn through the origin perpendicular to the tangent planes to the cone

$$ax^2 + by^2 + cz^2 + 2a'yz + 2b'zx + 2c'xy = 0;$$

shew that they will generate the cone which has for its equation

$$(bc - a'^2) x^2 + (ca - b'^2) y^2 + (ab - c'^2) z^2 \\ + 2(b'c' - aa') yz + 2(c'a' - bb') zx + 2(a'b' - cc') xy = 0.$$

Shew also that if lines be drawn through the origin perpendicular to the tangent planes to the second cone they will generate the first cone.

III. *Surfaces in general.*

187. If a sphere pass through the origin of co-ordinates and its centre is on the surface defined by the equation

$$\{3(x^2 + y^2 + z^2) - b^2\}^2 = (x + y + z)^2(x^2 + y^2 + z^2),$$

the sum of the spherical surfaces cut off by the co-ordinate planes is constant and  $= 2\pi b^2$ .

188. A straight line drawn from the centre of an ellipsoid meets the ellipsoid in  $P$  and the sphere on the diameter  $2a$  in  $Q$ ; shew that the tangent planes at  $P$  and  $Q$  contain a constant angle  $\alpha$  if the co-ordinates of  $P$  satisfy the equation

$$(x^2 + y^2 + z^2) \left( \frac{x^2}{a^4} + \frac{y^2}{b^4} + \frac{z^2}{c^4} \right) = \sec^2 \alpha.$$

189. If a straight line be drawn from the centre of an hyperboloid whose equation is  $\frac{x^2}{a^2} + \frac{y^2}{b^2} - \frac{z^2}{c^2} = 1$  to meet the surface in  $P, P'$ , and a point  $Q$  be taken in  $CP$  produced such that  $CP^2 = l(QP + QP')$ , where  $l$  is a constant, shew that the locus of  $Q$  is defined by the equation

$$\left( \frac{x^2}{a^2} + \frac{y^2}{b^2} - \frac{z^2}{c^2} \right)^2 = \frac{x^2 + y^2 + z^2}{4l^2}.$$

190. The shadow of a given ellipsoid thrown by a luminous point on the plane which passes through two of the principal axes has its centre on the curve in which the same plane intersects the ellipsoid; shew that the equation to the locus of the luminous point is

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = \left( \frac{z^2}{c^2} - 1 \right)^2.$$

191. If light fall from a luminous point whose co-ordinates are  $\alpha, \beta, \gamma$ , on a surface whose equation is  $xyz = m^3$ , the boundary of light and shade lies on an hyperboloid of one or two sheets according as the product of  $\alpha, \beta, \gamma$ , is negative or positive.

192. Tangent planes to an ellipsoid are drawn at a given distance from the centre; find the projections on the principal planes of the curve which is the locus of the points of contact. In what case will one of these projections consist of two straight lines?

193. Shew that the surfaces whose equations are

$$(x^2 + y^2)(a^2 - z^2) - c^2y^2 = 0,$$

$$\text{and } (x - c)^2 + y^2 + z^2 - a^2 = 0,$$

touch one another; and that the projection of the curve of contact on one of the co-ordinate planes is a circle, and on the other a parabola.

194. Two circles have a common diameter  $AB$  and their planes are inclined to each other at a given angle; on  $PP'$  any chord of one of them parallel to  $AB$  is described a circle with its plane parallel to that of the other circle; shew that the surface generated by these circles is an ellipsoid the squares of whose axes are in arithmetical progression.

195. Find the equation to the surface generated by straight lines drawn through the origin parallel to normals to the surface  $\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$  at points where it is intersected by the surface  $\frac{x^2}{a^2 - k^2} + \frac{y^2}{b^2 - k^2} + \frac{z^2}{c^2 - k^2} = 1$ ;  $k$  being greater than  $c$  and less than  $a$ .

196. An ellipse of given excentricity moving with its plane parallel to the plane of  $(y, z)$  and touching at the extremities of its axes the planes of  $(x, y)$  and  $(x, z)$  always passes through the curve whose equations are  $y = x$ ,  $cz = x^2$ ; find the equation to the surface generated, and determine the volume bounded by the surface and two given positions of the generating ellipse.

197. Shew that the surface determined by

$$z = F\left(\frac{ax + by + cz}{a'x + b'y + c'z}\right)$$

is cut by planes parallel to the plane of  $(x, y)$  in straight lines.

198. A tangent line to the surface of the ellipsoid

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$$

passes through the axis of  $z$ , and a given curve in the plane of  $(x, y)$ ; shew how to find the equation to the surface generated by it. Ex. The curve being  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = m^2$ , shew that the surface is a cone of the second order.

199. Find the differential equation to the surface generated by a straight line which passes through two given curves and remains parallel to the plane of  $(x, y)$ . Shew that the equation  $x = (y - b) \tan \frac{z}{a}$  represents such a surface.

200. Determine the surface generated by a straight line which moves parallel to the plane of  $(x, y)$ , and passes through the axis of  $z$  and through the curve given by

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1, \quad \frac{x}{a} + \frac{y}{b} = 1.$$

201. Determine the surface generated by a straight line which moves parallel to the plane of  $(x, y)$  and passes through the following curves (1) and (2):

$$\frac{x^2}{a^2} + \frac{z^2}{c^2} = 1, \quad y = 0 \dots \dots \dots (1),$$

$$\frac{y^2}{b^2} + \frac{z^2}{c^2} = 1, \quad x = 0 \dots \dots \dots (2).$$

202. A surface is generated by a straight line which always passes through the two fixed straight lines

$$y = mx, \quad z = c; \quad \text{and} \quad y = -mx, \quad z = -c;$$

prove that the equation to the surface generated is of the form

$$\frac{mcx - yz}{c^2 - z^2} = f\left(\frac{mzx - cy}{c^2 - z^2}\right).$$

203. If a ruled surface be such that at any point of it a straight line can be drawn lying wholly on the surface and intersecting the axis of  $z$ , then at every point of the surface

$$x^2 \frac{d^2 z}{dx^2} + 2xy \frac{d^2 z}{dx dy} + y^2 \frac{d^2 z}{dy^2} = 0.$$

204. The general equation to surfaces generated by a straight line which is always parallel to the plane

$$lx + my + nz = 0,$$

$$\begin{aligned} \text{is } \left(m + n \frac{dz}{dy}\right)^2 \frac{d^2 z}{dx^2} - 2 \left(m + n \frac{dz}{dy}\right) \left(l + n \frac{dz}{dx}\right) \frac{d^2 z}{dx dy} \\ + \left(l + n \frac{dz}{dx}\right)^2 \frac{d^2 z}{dy^2} = 0. \end{aligned}$$

205. Find the equation to the surface generated by a line which always passes through each of two given lines in space, and also through the circumference of a circle whose plane is parallel to them both, and whose centre bisects the shortest distance between them.

206. A surface is generated by a straight line which always intersects the line  $\frac{x-a}{l} = \frac{y-b}{m} = \frac{z-c}{n}$ , and is parallel to the plane  $\lambda x + \mu y + \nu z = 0$ ; find the functional equation to the surface; also find the differential equation.

207. Find the general functional equation to surfaces generated by the motion of a straight line which always intersects and is perpendicular to a given straight line.

If a surface whose equation referred to rectangular axes is

$$ax^2 + by^2 + cz^2 + 2a'yz + 2b'zx + 2c'xy + 2a''x + 2b''y + 2c''z + 1 = 0$$

be capable of generation in this manner, shew that

$$a + b + c = 0, \quad aa'' + bb'' + cc'' = 2a'b'c' + abc.$$

208. Shew that  $xyz = c(x^2 - y^2)$  represents a conoidal surface.

209. Shew that the condition in order that

$$ax^2 + by^2 + cz^2 + 2a''x + 2b''y + 2c''z = f$$

may represent a conical surface is

$$\frac{a''^2}{a} + \frac{b''^2}{b} + \frac{c''^2}{c} + f = 0.$$

210. Shew that in order that the equation

$$ax^2 + by^2 + cz^2 + 2a'yz + 2b'zx + 2c'xy + 2a''x + 2b''y + 2c''z + f = 0$$

may represent a cylindrical surface, it is necessary that

$$\frac{a''}{aa' - b'c'} + \frac{b''}{bb' - c'a'} + \frac{c''}{cc' - a'b'} = 0.$$

Is this condition sufficient as well as necessary?

211. Explain the two methods of generating a developable surface; find the differential equation to developable surfaces from each mode of generation.

212. Are the following surfaces developable?

$$(1) \quad xyz = a^3; \quad (2) \quad z - c = \sqrt{xy}.$$

213. Find the differential equation to a surface whose tangent plane at any point includes with the three co-ordinate planes a pyramid of constant volume; shew that the surface is generally developable.

214. Find the equation to the surface in which the tangent plane at  $(x, y, z)$  meets the axis of  $z$  at a distance from the origin equal to that of  $(x, y, z)$  from the origin.

215. Find the equation to the surface in which the tangent plane at  $(x, y, z)$  meets the axis of  $z$  at a distance  $\frac{k^{n+1}}{z^n}$  from the origin. If  $n = 1$ , give that form to the arbitrary function which will produce the equation to an ellipsoid.

216. A plane passes through  $(0, 0, c)$  and touches the line

$$x^2 + y^2 = a^2, \quad z = 0;$$

determine the locus of the ultimate intersections of the plane.

217. Three points move with given uniform velocities along three rectangular axes from given positions; shew how to find the surface to which the plane passing through their contemporaneous positions is always a tangent.

218. Spheres of constant radius  $r$  are described passing through the origin; find the envelope of the planes of contact of tangent cones having a fixed vertex at the point  $(a, b, c)$ .

219. Find the locus of the ultimate intersections of a series of planes touching two parabolas which lie in planes perpendicular to each other and have a common vertex and axis.

220. The sphere  $x^2 + y^2 + z^2 = 2ax + 2by + 2cz$  is cut by another sphere which passes through the origin and has its centre on the surface

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1;$$

shew that the equation to the envelope of the planes of intersection is

$$\frac{1}{ax} + \frac{1}{by} + \frac{1}{cz} = 0.$$

221. From each point of the exterior of two concentric ellipsoids, whose axes are in the same directions, tangent planes are drawn to the surface of the interior ellipsoid; shew that all the planes of contact corresponding to the several points of the exterior surface touch another concentric ellipsoid.

222. From any point  $P$  in the surface of an ellipsoid straight lines are drawn so as each to pass through one of three conjugate diameters, and be parallel to the plane containing the other two; these lines meet the surface again in  $P_1, P_2, P_3$ ; find the equation to the plane which passes through these points, and the locus of the ultimate intersections of all such planes, the diameters remaining fixed while  $P$  moves; and shew that its volume =  $\frac{3}{35}$  of that of the ellipsoid.



223. A cone envelopes the ellipsoid

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1,$$

and its vertex moves on the similar ellipsoid

$$\frac{(x - \alpha)^2}{a^2} + \frac{(y - \beta)^2}{b^2} + \frac{(z - \gamma)^2}{c^2} = 1;$$

shew that the plane of contact touches the surface

$$\left(\frac{\alpha x}{a^2} + \frac{\beta y}{b^2} + \frac{\gamma z}{c^2} - 1\right)^2 = n^2 \left(\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2}\right).$$

224. Three ellipsoids are represented by the following equations :

$$Ax^2 + By^2 + Cz^2 = a^2 \dots\dots\dots (1),$$

$$A^2x^2 + B^2y^2 + C^2z^2 = a^2 \dots\dots\dots (2),$$

$$A^3x^2 + B^3y^2 + C^3z^2 = a^2 \dots\dots\dots (3).$$

If tangent planes be drawn to (2) from every point of (1), the successive positions of the planes of contact will touch (3).

225. One axis of an ellipsoid is gradually increased while the volume remains constant, and the section containing the other axes similar; find the enveloping surface.

226. Prove that an infinite number of plane centric sections of an hyperboloid of one sheet may be drawn, each possessing the following property, viz. that the normals to the surface at the curve of section all pass through one or other of two straight lines lying in the same plane with the two possible axes.

Shew that these centric planes envelope the asymptotic cone.

227. Prove that the envelope of a sphere of which any one of one series of circular sections of an ellipsoid is a diametral plane, is another ellipsoid touching a sphere described on the mean axis of the former ellipsoid as diameter, in a plane perpendicular to any one of the same series of circular sections.

228. Find the envelope of a sphere of constant radius having its centre on a given circle, and determine the section by a tangent plane perpendicular to the plane of the circle.

229. From every point of the surface

$$4z (lx^2 + ly^2) + W (x^2 + y^2) = 0 \dots\dots\dots(1),$$

as vertex enveloping cones are drawn to the paraboloid

$$Wz - (lx^2 + ly^2) = 0 \dots\dots\dots(2).$$

Shew that the planes of contact all touch the surface

$$4z (lx^2 + ly^2) - (l^2x^2 + l^2y^2) = 0.$$

Also shew that the surface (1) contains the directrices of all the sections of (2) made by planes through its axis.

IV. *Curves.*

230. A curve is determined by the equations  $x + z = a$ , and  $x^2 + y^2 + z^2 = a^2$ ; find the points where the tangent makes an angle  $\beta$  with the axis of  $z$ .

231. Find the equations to the tangent to the curve

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} - \frac{z^2}{c^2} = 0, \quad \frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1.$$

232. Find the equations which determine the tangent to the curve

$$y^2 = ax - x^2, \quad z^2 = a^2 - ax.$$

233. Find the equations to the curve traced out on the surface

$$\frac{x^2}{a} + \frac{y^2}{b} = z$$

by the extremities of the latera recta of sections made by planes passing through the axis of  $z$ .

234. Having given the equations to a curve in space referred to three rectangular axes, find the length of the perpendicular from the origin upon the tangent at any point.

$$\text{Ex. } x = a \cos \theta, \quad y = a \sin \theta, \quad z = \frac{a}{2} (e^\theta + e^{-\theta}).$$

Prove that if the perpendicular be invariable either the curve or its involute lies on the surface of a sphere.

235. A perpendicular is drawn from the origin on the tangent to a curve at the point  $(x, y, z)$ . If  $x', y', z'$  are the co-ordinates of the foot of the perpendicular, shew that

$$x' = x - \frac{dx}{ds} \left( x \frac{dx}{ds} + y \frac{dy}{ds} + z \frac{dz}{ds} \right),$$

$$y' = y - \frac{dy}{ds} \left( x \frac{dx}{ds} + y \frac{dy}{ds} + z \frac{dz}{ds} \right),$$

$$z' = z - \frac{dz}{ds} \left( x \frac{dx}{ds} + y \frac{dy}{ds} + z \frac{dz}{ds} \right).$$

236. Shew how to determine the locus of the feet of the perpendiculars from the origin upon the tangent lines to a given curve.

The equations to the curve being  $x = a \cos \theta$ ,  $y = a \sin \theta$ ,  $z = c\theta$ , shew that the locus is a curve lying on the surface

$$\frac{x^2 + y^2}{c^2} - \frac{z^2}{a^2} = \frac{a^2}{c^2}.$$

237. A curve is traced on a sphere so that its tangent makes always a constant angle with a fixed plane; find its length from cusp to cusp.

238. Find the equation to the normal plane to a curve at any proposed point; if the normal plane always passes through a fixed straight line, shew that the curve is a circle.

239. Find the direction cosines of the line in which the normal plane at any point of a curve meets the osculating plane at that point.

240. Find the equation to the osculating plane at any point of the curve

$$x = a \cos t, \quad y = b \sin t, \quad z = ct.$$

241. If  $(x, y, z)$  be any point on a curve, shew what plane is represented by the equation

$$(x' - x) \frac{d^2x}{ds^2} + (y' - y) \frac{d^2y}{ds^2} + (z' - z) \frac{d^2z}{ds^2} = 1.$$

242. Find the value of the radius of absolute curvature from the consideration that it is the perpendicular from  $(x, y, z)$  on the plane in the preceding example.

243. Find the radius of curvature of the curve

$$\frac{x}{h} + \frac{y}{k} = 1, \quad x^2 + z^2 = a^2,$$

at any point.

244. Determine the radii of absolute and spherical curvature and the co-ordinates of the centres in the case of a helix.

245. Required the locus from which three given spheres will appear of equal magnitudes.

246. A triangular pyramid upon a given base is such that given lengths being measured along the three edges from the base the remainders of the edges are always equal to one another; shew that the locus of the vertex is a conic section.

247. Required the locus of the vertex of a cone of revolution whose base is a given conic section.

248. A point moves on the cone  $y^2 + z^2 = m^2x^2$  so that the tangent to its path is inclined to the axis of the cone at a constant angle  $\beta$ ; shew that the locus of the point is determined by the equation to the cone together with the equation

$$k \log \frac{x}{c} = \sin^{-1} \frac{y}{mx},$$

where  $k = \frac{\sqrt{(\tan^2\beta - m^2)}}{m}$ , and  $c$  is a constant.

249. Shew that a curve drawn on the surface of a right cone so as to cut the generating lines at a constant angle is a curve of constant inclination to the base of the cone, and that the consecutive distances between the points in which it cuts any one generator are in geometrical progression.

250. From the vertex of a right cone two curves are drawn on the surface cutting all the generating lines at constant angles which are complementary; shew that the sum of the inverse squares of the arcs intercepted between the vertex and a given circular section is independent of the magnitudes of the complementary angles.

251. Find the curves of greatest inclination to the co-ordinate planes on the surface of which the equation is  $xyz = a^3$ .

252. Find the curves of greatest inclination to the plane of  $(x, y)$  on the surface  $cz = xy$ .

253. Tangent planes are drawn to the surface  $cz = xy$  at all points where this surface is intersected by the cylinder  $x^2 = ay$ ; find the equations to the edge of regression of the developable surface formed by the intersection of these tangent planes.

254. Find the angle between the osculating planes at two consecutive points of a curve.

255. The shortest distance between the tangents at two consecutive points of a curve of double curvature is  $\frac{\delta\phi(\delta s)^2}{4\rho}$ , where  $\delta s$  is the length of the arc between the two points,  $\delta\phi$  is the angle between the osculating planes at the two points, and  $\rho$  is the radius of absolute curvature.

256. Shew that the edge of the developable surface formed by the locus of the ultimate intersections of normal planes of a curve of double curvature is the locus of the centres of spherical curvature of the curve.

Find the locus of the centres of spherical curvature of a helix.

257. Find the equation to the surface on which are found all the evolutes of the curve  $x^2 = az$ ,  $x = y$ .

258. Shew that the tangent lines to the curve represented by the equations

$$x^2 + y^2 + z^2 = a^2;$$

$\sqrt{x} + \sqrt{y} + \sqrt{z} = c$  are cut perpendicularly by each of the series of surfaces

$$x^{\frac{1}{2}} + (\mu - 1)y^{\frac{1}{2}} = \mu z^{\frac{1}{2}},$$

$\mu$  being an arbitrary parameter.

259. A curve is traced on a surface; shew that the radius of absolute curvature at any point of this curve is the same as the radius of curvature of the section of the surface made by the osculating plane of the curve at that point.

260. A curve is traced on a sphere; shew that generally the radius of the sphere is the radius of spherical curvature of the curve; but that this does not hold if the curve be a *plane* curve, or plane for an indefinitely short period at any point.

261. If the normal plane of a curve constantly touches a given sphere the curve is rectifiable.

262. If  $(x', y', z')$  be the point in the locus of the centres of curvature corresponding to the point  $(x, y, z)$  on a curve,  $\rho$  the radius of curvature at  $(x, y, z)$ , and  $s'$  the arc of the above locus, shew that  $\frac{d\rho}{ds}$  is the cosine of the angle between the tangent at  $(x', y', z')$  and the direction of  $\rho$ .

263.  $P$  is any point in a curve, and from the pole  $S$  the line  $SQ$  is drawn parallel to the tangent at  $P$  and of such a length that the parallelogram of which  $SP, SQ$  are sides has an invariable area; shew that the tangent to the locus of  $Q$  is parallel to  $SP$ .

V. *Curvature of Surfaces.*

264. Determine the conditions necessary in order that the surfaces whose equations are

$$ax^2 + by^2 + cz^2 + 2a'yz + 2b'zx + 2c'xy + 2z = 0,$$

$$Ax^2 + By^2 + Cz^2 + 2A'yz + 2B'zx + 2C'xy + 2z = 0,$$

may have their principal radii of curvature at the origin equal; and shew that if these conditions be fulfilled any sections of the two surfaces parallel to the plane of  $(x, y)$  will be similar.

265. Obtain the quadratic equation for determining the principal radii of curvature at any point of the surface

$$\phi(x) + \chi(y) + \psi(z) = 0;$$

and find the condition that the principal curvatures may be equal and opposite.

266. The locus of the points on the hyperboloid

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} - \frac{z^2}{c^2} = 1,$$

for which the principal curvatures are equal and opposite, has for its projection on the plane of  $(x, y)$  the ellipse

$$\frac{x^2}{a^2} (a^2 + c^2) + \frac{y^2}{b^2} (b^2 + c^2) = a^2 + b^2.$$

267. Shew that the principal radii of curvature are equal in magnitude and opposite in sign at every point of the surface determined by

$$z = (y - b) \tan \frac{x}{a}.$$

268. The trace on the plane of  $(y, z)$  of the locus of the extremities of the principal radii of curvature of the ellipsoid whose equation is

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$$



is given by the equation

$$\{(by)^{\frac{2}{3}} + (cz)^{\frac{2}{3}} - (b^2 - c^2)^{\frac{2}{3}}\} \left\{ \frac{b^2 y^2}{(a^2 - b^2)^2} + \frac{c^2 z^2}{(a^2 - c^2)^2} - 1 \right\} = 0.$$

269. Among the surfaces included in the equation

$$x \frac{dz}{dx} + y \frac{dz}{dy} = 0,$$

find that in which the principal radii of curvature are equal but of opposite sign.

270. Find the surface of revolution at every point of which the radii of curvature are equal in magnitude and opposite in sign.

271. If  $a$ ,  $b$  be the principal radii of curvature at any point of a surface referred to the tangent plane at that point as the plane of  $(x, y)$  and the principal planes as planes of  $(x, z)$  and  $(y, z)$ , then will the locus of the circles of curvature of all sections of the surface at the origin be

$$(x^2 + y^2 + z^2) \left( \frac{x^2}{a} + \frac{y^2}{b} \right) = 2z(x^2 + y^2).$$

272. Find the radius of curvature in any normal section of the surface

$$Ax^2 + By^2 + Cz^2 + 2A'yz + 2B'zx + 2C'xy + Ez = 0,$$

at the origin; and shew that the sum of the reciprocals of the radii of curvature in sections at right angles to each other is constant.

273. Required the sum of the principal radii of curvature at any point of a curved surface in terms of the co-ordinates of that point.

The equation to the surface being  $f(x, y, z) = 0$ , express the result by partial differential coefficients of  $f(x, y, z)$ .

274. If  $r = f(\theta, \phi)$  be the equation to a surface referred to the tangent plane at the origin as the plane of  $(x, y)$ , then

the radius of curvature at the origin of a normal section inclined at any angle  $\phi$  to the plane of  $(x, z)$  is

$$= -\frac{1}{2} \text{ limit of } \frac{dr}{d\theta} \text{ when } \theta = \frac{\pi}{2}.$$

Ex. In the surface  $xy = az$ , shew that  $\rho = \frac{a}{\sin 2\phi}$ .

275. If the surface  $(x - a)^2 + (y - b)^2 = (z - c)^2$  have contact of the second order with the surface  $z = f(x, y)$ , shew that the relations

$$\frac{p^2 - 1}{r} = \frac{q^2 - 1}{t} = \frac{pq}{s}$$

must be satisfied at the point of contact.

276. At the point in which the surface

$$\left(\frac{x^2}{a^2} - 1\right)^2 + \left(\frac{y^2}{b^2} - 1\right)^2 = \frac{z^2}{c^2} + 1$$

meets the axis of  $z$ , an elliptic paraboloid may be found, which has at its vertex a complete contact of the third order with the surface.

277. Find the radius of curvature of a normal section of an ellipsoid made by a plane inclined to the meridian at any given angle.

278. Shew that the locus of the focus of an ellipse rolling along a straight line is a curve such that if it revolves about that line, the sum of the curvatures of any two normal sections at right angles to one another will be the same for all points of the surface generated.

279. In any surface of the second order the tangents to the lines of curvature at any point are parallel to the axes of any plane section parallel to the tangent plane to the surface at that point.

280. Obtain the differential equation to the projection on one of the co-ordinate planes of the lines of curvature of a

surface. Apply the equation to determine the lines of curvature of a surface of revolution.

281. Determine the lines of curvature on the surface  $xy = az$ .

282. Find the radius of curvature of any normal section of a surface at a given point.

If  $y' - y = m(x' - x)$  be the projection on the plane of  $(x, y)$  of the tangent line to the curve of section, shew that the values of  $m$  corresponding to the principal sections of the surface  $\frac{y}{z} = f\left(\frac{x}{z}\right)$  at the point  $(x, y, z)$  are

$$\frac{y}{z} \text{ and } -\frac{x + pz}{y + qz}.$$

283. The links of a chain are circular, being of the form of the surface generated by the revolution of a circle whose radius is one inch about a line in its own plane at a distance of four inches from the centre; apply Euler's formula for the curvature of surfaces to shew that if one link be fixed, the next cannot be twisted through an angle greater than  $60^\circ$  without shortening the chain.

284. The normals about any point of a curve surface ultimately pass through two fixed straight lines in the planes of greatest and least curvature.

285. An annular surface is generated by the revolution of a circle about an axis in its own plane; prove that one of the principal radii of curvature at any point of the surface varies as the ratio of the distance of this point from the axis to its distance from the cylindrical surface described about the axis and passing through the centre of the circle.

286. If  $\rho, \rho'$  be the greatest and least radii of curvature of a curve surface at a given point,  $\phi, \psi$  the angles which the normal to the surface at the given point makes with the axes of  $x$  and  $y$ , shew that

$$\frac{1}{\rho} + \frac{1}{\rho'} = \frac{d}{dx} \cos \phi + \frac{d}{dy} \cos \psi.$$

287. Define an umbilicus. In what sense do you understand that there is an infinite number of lines of curvature at an umbilicus? And from this consideration deduce the partial differential equations which exist at such points.

288. Find the umbilici of the surface

$$2z = \frac{x^2}{\alpha} + \frac{y^2}{\beta}.$$

289. Find the umbilici of the surface  $xyz = a^3$ .

290. Prove that the radius of normal curvature of the surface  $xyz = a^3$  at an umbilicus is equal to the distance of the umbilicus from the origin of co-ordinates.

291. A sphere described from the origin with radius  $\frac{abc}{ac + ab + bc}$  will touch the surface

$$\left(\frac{x}{a}\right)^{\frac{2}{3}} + \left(\frac{y}{b}\right)^{\frac{2}{3}} + \left(\frac{z}{c}\right)^{\frac{2}{3}} = 1$$

in points which are umbilici.

292. Shew that a sphere whose centre is at the origin and whose radius  $r$  is determined by the equation

$$r^{\frac{2n}{n-2}} = a^{\frac{2n}{n-2}} + b^{\frac{2n}{n-2}} + c^{\frac{2n}{n-2}}$$

will touch the surface

$$\left(\frac{x}{a}\right)^n + \left(\frac{y}{b}\right)^n + \left(\frac{z}{c}\right)^n = 1$$

at umbilici; and the radius of normal curvature of the surface at these points is  $\frac{r}{n-1}$ .

293. Determine whether there is an umbilicus on the surface

$$(x^2 + y^2 + z^2)^2 = 4a^2(x^2 + y^2).$$

294. If  $R$  be the radius of absolute curvature at any point of a curve defined by the intersection of two surfaces  $u_1 = 0$ ,  $u_2 = 0$ , and  $r_1, r_2$  be the radii of curvature of the sections of  $u_1 = 0, u_2 = 0$ , made by the tangent planes to  $u_2 = 0, u_1 = 0$ , respectively at that point, prove that  $R, r_1, r_2$  will be connected by the relation

$$\frac{1}{R^2} = \frac{1}{r_1^2} - \frac{2 \cos \theta}{r_1 r_2} + \frac{1}{r_2^2},$$

$\theta$  being the angle between the tangent planes

295. Two surfaces touch each other at the point  $P$ ; if the principal curvatures of the first surface at  $P$  be denoted by  $a \pm b$ , those of the second by  $a' \pm b'$ , and if  $\varpi$  be the angle between the principal planes to which  $a + b, a' + b'$ , refer, and  $\delta$  the angle between the two branches at  $P$  of the curve of intersection of the surfaces, shew that

$$\cos^2 \delta = \frac{a^2 + a'^2 - 2aa'}{b^2 + b'^2 - 2bb' \cos 2\varpi}.$$

296. Prove that at any point of a developable surface, the curvature of any normal section varies as the square of the sine of the angle which this section makes with the generating line, and that at different points along the same generating line the principal radius of curvature varies as the distance from the point of intersection of consecutive generating lines.

297. A surface is generated by the motion of a straight line which always intersects a fixed axis. If  $P$  be any point in this axis at a distance  $x$  from the origin,  $\phi$  the angle which the generating line through this point makes with the axis, and  $\psi$  the angle which the plane through the axis and the generating line makes with its initial position, prove that the principal radii of curvature at  $P$  are

$$\cot \frac{\phi}{2} \frac{dx}{d\psi} \quad \text{and} \quad -\tan \frac{\phi}{2} \frac{dx}{d\psi}.$$

298. The shortest distance between two points on a curved surface never coincides with a line of curvature unless it be a plane curve.

VI. *Miscellaneous Examples.*

299. Find the condition that must hold in order that the equations

$$ax + c'y + b'z = 0, \quad by + a'z + c'x = 0, \quad cz + b'x + a'y = 0,$$

may represent a straight line; and shew that in that case the straight line is determined by the following equations:

$$x(aa' - b'c') = y(bb' - c'a') = z(cc' - a'b').$$

300. Shew that the six planes which bisect the interior angles of a tetrahedron meet in a point.

301. Shew that the three planes which bisect the exterior angles round one face of a tetrahedron and the three planes which bisect the interior angles formed by the other three faces meet in a point.

302. If  $\alpha = 0$ ,  $\beta = 0$ ,  $\gamma = 0$ ,  $\delta = 0$  be the equations to the faces of a tetrahedron expressed in a suitable form,  $A$ ,  $B$ ,  $C$ ,  $D$  the areas of the respective faces, shew that

$$A\alpha + B\beta + C\gamma + D\delta = \text{a constant.}$$

Interpret  $A\alpha + B\beta + C\gamma = 0$ .

303. If  $\alpha = 0$ ,  $\beta = 0$ ,  $\gamma = 0$  be the equations to three planes which form a trihedral angle, the equation to a cone of the second degree which has its vertex at the angular point and touches two of the planes at their intersections with the third, is  $\gamma^2 - k\alpha\beta = 0$ .

304. Give the geometrical interpretation of the equation  $uu' = kvv'$ , where  $k$  is a constant, and the other letters denote linear functions of  $x$ ,  $y$ ,  $z$ . Hence shew that there must exist surfaces of the second order which contain straight lines.

305. If  $u_1 = 0$ ,  $u_2 = 0$ ,  $u_3 = 0$  be the equations to three planes, interpret the equation

$$\sqrt{\left(\frac{u_1}{a}\right)} + \sqrt{\left(\frac{u_2}{b}\right)} + \sqrt{\left(\frac{u_3}{c}\right)} = 0.$$

306. Also interpret  $Au_2u_3 + Bu_1u_3 + Cu_1u_2 = 0$ .

307. If  $u, v, w$  are linear functions of  $x, y, z$  shew that  $uv = w^2$  represents a conical surface; and shew that the equation to the tangent plane is  $\lambda^2u - 2\lambda w + v = 0$ .

308. Let  $v = 0$  be the equation to a surface of the second order;  $u_1 = 0$  and  $u_2 = 0$  the equations to two planes; shew that by giving a suitable value to the constant  $\lambda$  the equation  $v + \lambda u_1u_2 = 0$  will represent any surface of the second order which passes through the intersections of the two planes with the given surface.

309. If  $t = 0, u = 0, v = 0, w = 0$  be the equations to four given planes, and  $\lambda, \mu,$  be two arbitrary constants, shew that  $t + \lambda u = 0$  represents a plane which passes through a fixed straight line,  $t + \lambda u + \mu v = 0$  a plane which passes through a fixed point, and  $tw + \mu uv = 0$  a surface of the second order which contains four fixed straight lines.

310. If the equation to a surface of the second order be  $u_2 + 2u_1 + 1 = 0$ , where  $u_1$  and  $u_2$  represent the terms of the first and second order respectively, and tangent planes be drawn to the surface from any point of the plane determined by  $u_1 + 1 = 0$ , the planes of contact will all pass through the origin.

311. If  $A, B, C, D$  be the angular points of a tetrahedron;  $\alpha, \beta, \gamma, \delta$  the distances of any point  $P$  from the faces respectively opposite to them, prove that the general equation of the hyperboloid of one sheet of which  $AB, CD$  are generating lines is

$$l\alpha\gamma + m\alpha\delta + n\beta\gamma + p\beta\delta = 0.$$

312. If  $\alpha, \beta, \gamma, \delta$  have the same meaning as in the preceding example, shew that the general equation to a surface of the second order circumscribing the tetrahedron is

$$A\alpha\beta + B\alpha\gamma + C\alpha\delta + D\beta\gamma + E\beta\delta + F\gamma\delta = 0.$$

Determine the condition necessary in order that the straight

line  $\frac{\alpha}{l} = \frac{\beta}{m} = \frac{\gamma}{n}$  may touch the surface; and hence shew that the equation to the tangent plane at the point  $(\alpha = \beta = \gamma = 0)$  is

$$C\alpha + E\beta + F\gamma = 0.$$

313. Interpret  $u^2 + v^2 + w^2 = a$ , where  $u, v, w$  are linear functions of  $(x, y, z)$  and  $a$  is a constant.

314. Interpret  $u^2 + v^2 - w^2 = a$ .

315. Interpret  $u^2 + v^2 = a$ .

316. Interpret  $u^2 - v^2 = a$ .

317. Interpret  $u^2 = a$ .

318. Interpret  $u^2 + v^2 + w = 0$ .

319. Interpret  $u^2 - v^2 + w = 0$ .

320. Interpret  $u^2 + v = 0$ .

321. Interpret  $\theta = \text{constant}$ .

322. Interpret  $\phi = \text{constant}$ .

323. Interpret  $r = \text{constant}$ .

324. Interpret  $F(\theta) = 0$ .

325. Interpret  $F(\phi) = 0$ .

326. Interpret  $F(r) = 0$ .

327. Interpret  $F(r, \theta) = 0$ .

328. Interpret  $F(r, \phi) = 0$ .

329. Interpret  $F(\theta, \phi) = 0$ .

330. Describe the form of the surface represented by

$$\frac{\theta^2}{\alpha^2} + \frac{\phi^2}{\beta^2} = 1.$$

331. Interpret  $r \cos \theta = a$ .

332. Shew that the polar equation to a plane may be put in the form  $\frac{c}{r} = \cos \alpha \cos \theta + \sin \alpha \sin \theta \cos (\phi - \beta)$ .



333. If  $p$  be the perpendicular from the origin on the tangent plane at any point of the surface  $r = f(\theta, \phi)$

$$\frac{1}{p^2} = \frac{1}{r^2} + \frac{1}{r^4} \left( \frac{dr}{d\theta} \right)^2 + \frac{\operatorname{cosec}^2 \theta}{r^4} \left( \frac{dr}{d\phi} \right)^2.$$

334. Shew that the polar partial differential equation to conical surfaces with the origin as vertex is

$$\frac{d\phi}{dr} = 0, \quad \text{or} \quad \frac{d\theta}{dr} = 0.$$

335. Shew that the polar partial differential equation to surfaces of revolution about an axis perpendicular to the plane of reference is  $\frac{dr}{d\phi} = 0$ .

336. Shew that the polar partial differential equation to cylindrical surfaces perpendicular to the plane of reference is

$$\sin \theta \frac{dr}{d\theta} + r \cos \theta = 0.$$

337. Find the volume of a pyramid which has its vertex at the point  $(x, y, z)$ , and for its base the triangle formed by joining the points where the plane

$$\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$$

meets the co-ordinate axes.

338. Prove the formulæ for transforming from one set of rectangular axes to another. Shew that all six axes lie on a certain cone of the second order.

339. If two systems of rectangular axes have the same origin, and  $(a_1, b_1, c_1)$ ,  $(a_2, b_2, c_2)$ ,  $(a_3, b_3, c_3)$  be the direction cosines of one system with respect to the other, shew that

$$a_1^2 = (b_2c_3 - b_3c_2)^2, \quad a_2^2 = (b_3c_1 - b_1c_3)^2, \quad a_3^2 = (b_1c_2 - b_2c_1)^2.$$

340. Find the size of a cube which will stop up a tube of uniform bore, the section of which is a regular hexagon whose sides are given.

341. The stereographic projection of a cube on a plane perpendicular to its diagonal (the pole being in the diagonal produced) is an equilateral hexagon, whose angles are alternately greater and less. If the pole of projection be at a distance from the cube equal to its diagonal, the sines of two adjacent angles of the hexagon are as 8 to 5.

342. A cube being placed with one face in contact with a given plane, determine the position of a luminous point such that the shadow cast on the plane shall be an equilateral pentagon of which the diagonal of the above face is one side.

343. Find the locus of the centres of the sections of a surface of the second order made by planes which all pass through a fixed straight line.

344. Find the locus of the middle points of all chords drawn in a surface of the second degree, the length of each chord varying as the diameter parallel to it.

345. From different points of the straight line  $\frac{x}{a} = \frac{y}{b}$ ,  $z=0$ , asymptotic straight lines are drawn to the hyperboloid

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} - \frac{z^2}{c^2} = 1;$$

shew that they will all lie in the planes

$$\frac{x}{a} - \frac{y}{b} = \pm \frac{z}{c} \sqrt{2}.$$

346. Common tangent planes are drawn to the ellipsoids

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1, \text{ and } \frac{x^2}{a'^2} + \frac{y^2}{b'^2} + \frac{z^2}{c'^2} = 1;$$

shew that the perpendiculars upon them from the origin lie in the surface of the cone

$$(a^2 - a'^2)x^2 + (b^2 - b'^2)y^2 + (c^2 - c'^2)z^2 = 0.$$

347. Determine what must be the form of a column in order that it may appear to be of uniform thickness to an observer in a given position.

348. If the section of the surface whose equation is  $ax^2 + by^2 + cz^2 + 2a'yz + 2b'zx + 2c'xy + 2a''x + 2b''y + 2c''z + f = 0$ , by the plane whose equation is

$$lx + my + nz = p$$

be a rectangular hyperbola, then will

$$l^2(b+c) + m^2(c+a) + n^2(a+b) = 2a'mn + 2b'nl + 2c'lm.$$

349. If the section of the surface  $xy + yz + zx = a^2$  by the plane  $lx + my + nz = \delta$  be a parabola, then will

$$l^2 + m^2 + n^2 - 2mn - 2nl - 2lm = 0.$$

350. Shew that the asymptotes of any plane section of an hyperboloid are parallel to the straight lines in which the asymptotic cone is cut by the parallel central plane.

351. Two hyperboloids of one and two sheets respectively have the same asymptotic cone; shew that the sections of the hyperboloid of one sheet by tangent planes to the other hyperboloid will be ellipses of constant area if the points of contact lie on the curve of section of the surface by a certain concentric ellipsoid whose axes are in the ratio of the squares of the axes of the hyperboloids.

352. Through two straight lines given in space two planes are drawn so as to be at right angles; find the locus of their intersection.

353. A surface is generated by the intersection of planes cutting the axes of co-ordinates at distances from the origin equal to the respective co-ordinates of each point of the ellipsoid  $\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$ ; if the surface so generated be treated as the ellipsoid and the process repeated  $n$  times, the equation to the  $n^{\text{th}}$  surface will be

$$\left(\frac{x}{a}\right)^{\frac{2}{2n+1}} + \left(\frac{y}{b}\right)^{\frac{2}{2n+1}} + \left(\frac{z}{c}\right)^{\frac{2}{2n+1}} = 1.$$

354. A surface is generated by a variable circle whose plane is parallel to  $x + y = 0$ , and which always passes through the axes of  $x$  and  $y$  and the line  $y = x, z = c$ ; find the equation to the surface. Also find the volume between the origin and the plane  $x + y = c$ .

355. Two equal parabolas have their common vertex at the origin, their axes coincident with the axis of  $x$  in opposite directions, and their planes coincident with the planes of  $(x, y)$  and  $(x, z)$  respectively; a straight line is drawn intersecting these curves and parallel to the plane  $y = z$ ; find the locus of its trace on the plane of  $(y, z)$ .

356. Explain the nature of the surface defined by the equation

$$\frac{y}{b} = \left\{ \left( \frac{x^2}{a^2} + \frac{z^2}{c^2} \right)^{\frac{1}{2}} - 1 \right\} \left\{ \left( \frac{x^2}{a^2} + \frac{z^2}{c^2} \right)^{\frac{1}{2}} - m \right\}^{\frac{1}{2}}$$

near the points where it meets the hyperboloid

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$$

in the several cases in which  $m$  is  $>$  or  $<$  1.

357. Planes are drawn perpendicular to the tangent lines to the surface  $f(x, y, z) = 0$  at a point  $(x, y, z)$  in it; shew that if at that point

$$\frac{df}{dx} = 0, \quad \frac{df}{dy} = 0, \quad \frac{df}{dz} = 0.$$

$$\frac{d^2f}{dx^2} = u, \dots \frac{d^2f}{dy dz} = u', \dots$$

the locus of the ultimate intersections of the planes is the cone

$$(vw - u^2)(\xi - x)^2 + \dots + 2(v'w' - uu')(\eta - y)(\zeta - z) + \dots = 0.$$

358. Shew that the normal cone at a singular point of the surface

$$(x^2 + y^2 + z^2)(a^2x^2 + b^2y^2 + c^2z^2) - a^2(b^2 + c^2)x^2 - b^2(c^2 + a^2)y^2 - c^2(a^2 + b^2)z^2 + a^2b^2c^2 = 0$$

has for its equation

$$(b^2 - c^2) (\xi - x_0)^2 + (c^2 - a^2) \eta^2 + (a^2 - b^2) (\zeta - z_0)^2 - \frac{c^2 + a^2}{ca} \sqrt{(a^2 - b^2)} \sqrt{(b^2 - c^2)} (\zeta - z_0) (\xi - x_0) = 0,$$

$$\text{where } x_0 = \frac{c \sqrt{(a^2 - b^2)}}{\sqrt{(a^2 - c^2)}}, \quad y_0 = 0, \quad z_0 = \frac{a \sqrt{(b^2 - c^2)}}{\sqrt{(a^2 - c^2)}}$$

are the co-ordinates of the singular point.

359. If  $f(x, y, z, c) = 0$  represent a system of surfaces for which  $c$  is a variable parameter, shew how to find the locus of the points of contact of tangent planes to each one of the system, each tangent plane passing through a fixed point.

Ex.  $x^2 + y^2 = 2c(z - c)$ , each of the tangent planes passing through the origin.

360. If  $PN$  be the normal at the point  $P$  of any surface,  $a, b$  the principal radii of curvature at  $P$ ,  $r$  the radius of curvature of the normal section made by a plane inclined at an angle  $\theta$  to the principal section to which  $a$  refers,  $PQ = s$  an indefinitely small arc in this section, prove that if  $D$  be the minimum distance of the normals at  $P$  and  $Q$ ,  $c$  the distance from  $P$  of their point of nearest approach

$$D^2 = \frac{\sin^2 \theta \cos^2 \theta (a - b)^2 s^2}{b^2 \cos^2 \theta + a^2 \sin^2 \theta}, \quad c = \frac{ab (b \cos^2 \theta + a \sin^2 \theta)}{b^2 \cos^2 \theta + a^2 \sin^2 \theta}.$$

Express also  $\frac{D^2}{s^2}$  and  $c$  in terms of  $a, b$  and  $r$ .

361. A family of surfaces is defined by the equation

$$F\{x, y, z, \phi(a), \psi(a)\} = 0,$$

where  $a$  is a variable parameter and  $\phi(a)$  and  $\psi(a)$  arbitrary functions of  $a$ ; shew how to find the partial differential equation of the second order which belongs to the envelope, and prove that it assumes the form

$$rt - s^2 + Pr + Qs + Rt + S = 0,$$

where  $P, Q, R, S$  are functions of  $x, y, z, p$  and  $q$ .

362. If the equation to a system of surfaces contains an arbitrary parameter, shew that a curve can always be found which cuts them all at right angles; and if the equations to a system of curves contain two arbitrary parameters a surface cannot always be found which cuts them all at right angles.

363. Find the equations to a curve which cuts at right angles a series of ellipsoids which have their axes fixed in position and two of them of given length.

364. Prove that the section of a surface made by a plane parallel and indefinitely near to the tangent plane at any point in the immediate neighbourhood of that point is generally a conic section; and explain fully the peculiarity of the surface near a point where this conic section is a parabola.

365. Shew how to determine whether a curved surface has a tangent plane which touches it along a line. Examples:

$$(1) \quad (x^2 + y^2 + z^2)^2 = 4a^2 (x^2 + y^2).$$

$$(2) \quad b^2 z = (x^2 + y^2)^2 - a^2 (x^2 + y^2).$$

366. The extremities of the minor axis of the elliptic sections of a right cone made by parallel planes lie in two generating lines of the cone.

367. Shew how to cut from a given cone an hyperbola whose asymptotes shall contain the greatest possible angle.

368. What is the section of a right cone by a plane when the cutting plane is parallel to a generating line, but not perpendicular to the plane containing the axis and that line?

369. If two spheres exterior to each other be inscribed in a right cone touching it in two circles on the same side of the vertex, and a plane be drawn touching the spheres and cutting the cone; shew that the section is an ellipse, that the points of contact of the spheres with the plane are the foci, and that the planes of the two circles contain the directrices.

370. A conical surface is placed with its circular base in contact with a plane. It is then slit along a generating line

and the vertex pressed in a direction perpendicular to the plane, the base remaining in contact while the surface opens out; shew that the extremities of the separating edges trace hyperbolic spirals.

371. A triangle  $ABC$  revolves about a straight line which bisects the angle  $A$ , and the conical surface generated by the sides containing that angle is cut by a plane passing through the other side and perpendicular to the plane of the triangle in one of its positions; prove that if  $e$  be the excentricity of the section,  $e^2 = \frac{(b-c)^2}{a^2}$ .

372. If the cone  $x^2 + y^2 = z(mx + nz)$  cut any sphere which has its centre at the origin, shew that the projection on the plane of  $(x, z)$  of the curve of intersection is an hyperbola which has its centre at the origin.

373. A straight inelastic band is wrapped smoothly on the surface of a cone; shew that however long it may be, the two ends of either of its edges cannot be made to meet, if the vertical angle of the cone be greater than  $60^\circ$ .

374. Shew that an oblique cone on a circular base can be cut by a plane not parallel to the base, so that the section shall be a circle. Prove that the cone so cut off is similar to the whole cone.

375. If  $f(x, y, z) = 0$  be the equation to any surface which passes through the origin, and  $\phi(x, y, z)$  the sum of all the terms of lowest dimension in  $f(x, y, z)$ , shew that  $\phi(x, y, z) = 0$  is the locus of all the tangent lines at the origin.

376. Find the surface which touches at the origin the surface whose equation is

$$(x^2 + y^2 + z^2 + 2ax)^2 - 4(c^2 - a^2)(x^2 + z^2) = 0;$$

and shew that as  $a$  diminishes and ultimately vanishes, the tangent cone contracts and ultimately becomes a line, and that as  $a$  increases up to  $c$  it expands and finally becomes a plane.

377. A circle always touches the axis of  $z$  at the origin, and passes through a fixed straight line in the plane of  $(x, y)$ ; find the equation to the surface generated. Shew that the origin is a singular point, and that in its immediate neighbourhood the surface may be conceived to be generated by a circle having its plane parallel to that of  $(x, y)$ , and its radius proportional to  $z^2$ .

378. A circle revolves round a chord in its own plane, the direction cosines of which are  $l, m, n$ ; shew that the equation to the surface generated is

$$(lx + my + nz - h)^2 + [\sqrt{x^2 + y^2 + z^2} - (lx + my + nz)] - k]^2 = r,$$

where  $r$  is the radius of the circle;  $h$  and  $k$  the distances of the centre of the circle along and from the chordal line which is supposed to pass through the origin; determine the singular points of this surface and the equation to the tangent cone there; also determine the equation to the tangent planes which touch the surface along circles.

379. The traces of a surface of the second order on two planes at right angles to each other are parabolas with their axes parallel to the line of intersection of the planes; find the condition that the surface may be a developable surface; and prove that in that case the trace of the surface on a plane perpendicular to the first two will also be a parabola.

380. Find the locus of the point in which a perpendicular from the origin meets a plane which cuts off a constant volume from the co-ordinate planes.

381. An oblique cylinder stands on a great circle of a sphere; determine the curve of intersection of the sphere and cylinder, and find the area of the spherical surface included within the cylinder.

382. The locus of the intersection of generating lines of an hyperboloid of one sheet which pass through extremities of conjugate diameters of the smallest elliptic section is a similar ellipse parallel to it, whose axes are to the axes of the former as  $\sqrt{2}$  to 1.



383. Find the surface which is generated by a straight line which is always parallel to the plane of  $(x, y)$ , and passes through the axis of  $z$ , and also through the curve determined by  $xyz = a^3$ ,  $x^2 + y^2 = c^2$ .

384. A normal to the surface of which the equation is

$$x \cos nz - y \sin nz = 0,$$

moves along any one of its generating lines; determine the surface generated.

385. The equations to a system of lines in space, straight or curved, contain two arbitrary parameters; shew how to find whether the lines can be cut at right angles by a system of surfaces, and when they can shew how to find the equation to that system.

EXAMPLES. (1) Let the lines be a system of right lines, each of which intersects two given right lines which are perpendicular to each other but do not intersect. (2) Let the equations to the system of lines be  $Ax^a = By^b = Cz^c$ , where  $A, B, C$ , are arbitrary.

386. The equation to a system of surfaces contains one arbitrary parameter; normals are drawn at all points of one of these surfaces, and the lengths of the normals are taken proportional to the ultimate distances between the surface in question and a consecutive surface; shew how to find the equation to the locus of the extremities of the normals.

387. The equations to a system of right lines in space contain two arbitrary parameters; prove that when the roots of a certain quadratic are real and unequal, there are two planes passing through a given line of the system which contain consecutive lines.

388. A cavity is just large enough to allow of the complete revolution of a circular disk of radius  $c$ , whose centre describes a circle of the same radius  $c$ , while the plane of the disk is constantly parallel to a fixed plane and perpendicular

to that of the circle in which its centre moves. Shew that the volume of the cavity is  $\frac{2c^3}{3} (3\pi + 8)$ .

389. The solid of which the surface is determined by the equation  $\sqrt{x} + \sqrt{y} + \sqrt{z} = \sqrt{a}$  revolves round the fixed axis of  $z$  and makes for itself a cavity in a mass of yielding material; determine the form and magnitude of the cavity.

390. Shew that the surfaces represented by the equations

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1, \quad \frac{\alpha^2}{x^2} + \frac{\beta^2}{y^2} + \frac{\gamma^2}{z^2} = 1,$$

will touch each other in eight points if

$$\frac{\alpha}{a} + \frac{\beta}{b} + \frac{\gamma}{c} = 1;$$

and prove that if tangent planes be drawn at these points they will form a solid whose opposite faces are similar and parallel, and the volume of which is

$$\frac{4}{3} \frac{(abc)^{\frac{3}{2}}}{(\alpha\beta\gamma)^{\frac{3}{2}}}.$$

391. Normals are drawn to the ellipsoid

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$$

at every point of its curve of intersection with the sphere

$$x^2 + y^2 + z^2 = p^2;$$

prove that the equation to the curve in which the locus of these normals is cut by the plane of  $(y, z)$  is

$$\frac{b^2 y^2}{a^2 - b^2} + \frac{c^2 z^2}{a^2 - c^2} = a^2 - p^2.$$

392. Find the envelope of the plane

$$ax + \beta y + \gamma z = 1,$$

the parameters  $\alpha, \beta, \gamma$  being subject to the relations

$$a\alpha + b\beta + c\gamma = 1,$$

$$a^2\alpha^2 + b^2\beta^2 + c^2\gamma^2 = 1.$$

393. Determine the singular point on the surface

$$a^2x^2 + b^2y^2 = z^2(c - z),$$

and the locus of the tangent lines there.

394. The tangent plane to a surface  $S$  cuts an ellipsoid, and the locus of the vertex of the cone which touches the ellipsoid in the curve of intersection is another surface  $S'$ . Prove that  $S$  and  $S'$  are reciprocal; that is, that  $S$  may be generated from  $S'$  in the same manner as  $S'$  has been generated from  $S$ .

395. A plane moves so as always to cut off from an ellipsoid the same volume; shew that it will in every position touch a similar and concentric ellipsoid.

396. Prove that the tangent plane at any point of the surface

$$(ax)^2 + (by)^2 + (cz)^2 = 2(bcyz + caxx + abxy),$$

intersects the surface

$$ayz + bzx + cxy = 0$$

in two straight lines which are at right angles to one another.

397. A plane is drawn through the axis of  $y$  such that its trace upon the plane of  $(z, x)$  touches the two circles in which the plane of  $(z, x)$  meets the surface generated by the revolution round the axis of  $z$  of the circle

$$(x - a)^2 + z^2 = c^2,$$

where  $a$  is greater than  $c$ ; determine the curve of intersection of the plane and the surface.

398. Find the envelope of a sphere of constant radius having its centre on a given circle; and determine the section by a tangent plane perpendicular to the plane of the circle; when will this section be a lemniscate, and when will it consist of two circles?

399. A plane is drawn through a generating line of an hyperboloid of one sheet; shew that it meets the surface again in a straight line.

400. A plane moves so as always to enclose between itself and a given surface  $S$  a constant volume; prove that the envelope of the system of such planes is the same as the locus of the centres of gravity of the portions of the planes comprised within  $S$ .

401.  $OA, OB, OC$  are three straight lines mutually at right angles, and a luminous point is placed at  $C$ ; shew that when the quantity of light received upon the triangle  $AOB$  is constant, the curve which is always touched by  $AB$ , will be an hyperbola whose equation referred to the axes  $OA, OB$  is

$$(y - mx)(x - my) = mc^2,$$

where  $OC = c$ , and  $m$  is a constant quantity.

402. If  $u = f(x, y, z)$  be a rational function of  $x, y, z$ , and if  $u = 0$  be the equation to a surface, for a point  $(a, b, c)$  of which all the partial differential coefficients of  $u$  as far as those of the  $(n-1)^{\text{th}}$  order vanish, shew that the conical surface whose equation is

$$\left\{ (x-a) \frac{d}{da} + (y-b) \frac{d}{db} + (z-c) \frac{d}{dc} \right\}^n f(a, b, c) = 0$$

will touch the proposed surface at the point  $(a, b, c)$ .

403. Determine the condition to which the vertices of a system of cones which envelope an ellipsoid must be subject, in order that the centres of the ellipses of contact may be equidistant from the centre of the ellipsoid.

404. A plane passes through the vertex of the elliptic paraboloid  $2z = \frac{x^2}{a} + \frac{y^2}{b}$ ; shew that if the radius of curvature of all its sections of the surface at the vertex be equal to  $c$ , the normal to the plane at the origin will lie upon the surface

$$(x^2 + y^2)^2 = c^2 (x^2 + y^2 + z^2) \left( \frac{x^2}{b} + \frac{y^2}{a} \right).$$

405. Find the position in which an ellipsoid must be placed in order that its orthogonal projection may be circular, assuming that the plane on which it is projected must be parallel to its mean axis.

406. Find the equation to the surface generated by a straight line of given length, which moves parallel to the plane of  $(x, y)$ , with one end in the plane of  $(y, z)$  and the other on a given curve,  $x = \phi(z)$ , in the plane of  $(x, z)$ .

## RESULTS OF THE EXAMPLES.

1.  $\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 2.$

2.  $\frac{x}{a} + \frac{y}{b} - \frac{z}{c} = 1.$

3.  $\frac{y}{b} = \frac{z}{c}; \quad \frac{x}{a} = \frac{z}{c}.$

4.  $\frac{x}{a} = \frac{y}{b} = \frac{z}{c}.$

5.  $\frac{x}{a} = \frac{y-b}{-b} = \frac{z}{c}; \quad \frac{x-a}{-a} = \frac{y}{b} = \frac{z}{c}.$

6.  $\frac{2abc}{\sqrt{(a^2b^2 + b^2c^2 + c^2a^2)}}.$

7.  $\frac{2abc}{\sqrt{(a^2b^2 + b^2c^2 + c^2a^2)}}.$

8.  $\cos^{-1} \frac{ab}{\sqrt{(a^2 + c^2)} \sqrt{(b^2 + c^2)}}.$

9.  $\cos^{-1} \frac{3}{\sqrt{(a^2 + b^2 + c^2)} \sqrt{\left(\frac{1}{a^2} + \frac{1}{b^2} + \frac{1}{c^2}\right)}}.$

10.  $\cos^{-1} \frac{a^2 + b^2 - c^2}{a^2 + b^2 + c^2}.$

11.  $\frac{x}{a} = \frac{2y}{b} = \frac{2z}{c}.$

12. The equation may be written

$$(x \cos \beta - y \cos a)^2 + (y \cos \gamma - z \cos \beta)^2 + (z \cos a - x \cos \gamma)^2 = 0;$$

thus it represents the line

$$\frac{x}{\cos a} = \frac{y}{\cos \beta} = \frac{z}{\cos \gamma}.$$

13.  $30^\circ.$       14.  $\frac{Ax + By + Cz - D}{Aa + B\beta + C\gamma - D} = \frac{A'x + B'y + C'z - D'}{A'a + B'\beta + C'\gamma - D'}.$

15.  $D(Ax + By + Cz) = D(A'x + B'y + C'z);$  the condition is  
 $\left(\frac{D}{D'}\right)^2 = \frac{A^2 + B^2 + C^2}{A'^2 + B'^2 + C'^2}.$

16.  $(x - a)\{m(c - c') - n(b - b')\} + \&c. = 0.$

17. First obtain the equation to the plane which contains the two given lines ; this is

$$x(m_1n_2 - m_2n_1) + y(n_1l_2 - n_2l_1) + z(l_1m_2 - l_2m_1) = 0 ;$$

then find the equation to a plane which is perpendicular to the plane just determined and equally inclined to the two given lines.

21.  $\frac{5}{3}\sqrt{2}$ .

22.  $(x - a)(mm' - m'n) + \&c. = 0$ .

23. The question is indeterminate, since an unlimited number of lines can be drawn as required.

24. Suppose the given plane to be determined by

$$Ax + By + Cz = D ;$$

and suppose that the line of intersection is to lie in the plane of  $(x, y)$  ; then we may assume for the equation to the required plane  $Ax + By + \lambda z = D$ , and determine  $\lambda$  suitably.

27.  $60^\circ$ .

28. Four straight lines.

29. The condition is  $a^2 + b^2 + c^2 + 2abc = 1$ .

30.  $(p_1 \cos \alpha - p \cos \alpha_1)x + (p_1 \cos \beta - p \cos \beta_1)y + \lambda z = 0$ ,  
where  $(p_1 \cos \alpha - p \cos \alpha_1)^2 + (p_1 \cos \beta - p \cos \beta_1)^2 + \lambda(p_1 \cos \gamma - p \cos \gamma_1) = 0$ .

31. Let the given point be  $(a, b, c)$  and the equations to the given planes

$$Ax + By + Cz = D, \quad A'x + B'y + C'z = D' ;$$

the required equation is

$$(x - a)(BC' - B'C) + \&c. = 0.$$

37.  $\frac{p + p'}{\sqrt{(2 + 2k)}}$  and  $\frac{p - p'}{\sqrt{(2 - 2k)}}$ , where  $k = U + mm' + nn'$ .

41. Let  $r$  denote the distance between  $(\alpha, \beta, \gamma)$  and  $(x, y, z)$  ; then

$$\begin{aligned} (A^2 + B^2 + C^2)r^2 &= (A^2 + B^2 + C^2)\{(x - \alpha)^2 + (y - \beta)^2 + (z - \gamma)^2\} \\ &= \{B(z - \gamma) - C(y - \beta)\}^2 + \{C(x - \alpha) - A(z - \gamma)\}^2 + \{A(y - \beta) - B(x - \alpha)\}^2 \\ &\quad + \{A(x - \alpha) + B(y - \beta) + C(z - \gamma)\}^2 ; \end{aligned}$$

The last term is *constant* if  $(x, y, z)$  be in the given plane ; hence the *least* value of  $r^2$  is obtained by making the other three terms vanish.

44. The exceptional case is when the line of intersection of two of the planes is perpendicular to the third plane.

$$45. \quad \frac{1}{a} \left\{ x - \frac{b\gamma - c\beta}{a^2 + b^2 + c^2} \right\} = \frac{1}{b} \left\{ y - \frac{ca - a\gamma}{a^2 + b^2 + c^2} \right\} = \frac{1}{c} \left\{ z - \frac{a\beta - ba}{a^2 + b^2 + c^2} \right\}.$$

46. The condition is  $al + bm + cn = 0$ ; then the equations may be written

$$\frac{1}{l} \left\{ x - \frac{cm - bn}{l^2 + m^2 + n^2} \right\} = \&c. ; \text{ or thus, } \frac{1}{l} \left\{ x - \frac{cn - bm}{3mn} \right\} = \&c.$$

47. The equations to the perpendicular are

$$\frac{x}{mc - nb} = \frac{y}{na - lc} = \frac{z}{lb - ma}.$$

49. There are two such lines determined by the given equation  $lx + my + nz = 0$ , combined with

$$x \sqrt{(l^2 + n^2)} = \pm y \sqrt{(m^2 + n^2)}.$$

50. Let  $x, y, z$  be the co-ordinates of any point in the line; let  $l, m, n$  be proportional to its direction cosines; then it may be shewn that

$$\cos^2 a = \frac{(nx - lz)^2 (mx - ly)^2}{\{(nx - lz)^2 + (ny - mz)^2\} \{(mx - ly)^2 + (mz - ny)^2\}};$$

from this we may shew that  $\tan^2 a = A (mz - ny)^2$  where  $A$  is a symmetrical function of  $x, y, z, l, m, n$ .

$$52. \quad \frac{x}{a^2} = \frac{y}{b^2} = \frac{z}{c^2} = \frac{1}{\sqrt{(a^2 + b^2 + c^2)}}; \quad \frac{x}{a} = \frac{y}{b} = \frac{z}{c} = \frac{1}{\sqrt{3}}.$$

$$53. \quad \text{Here } e = \frac{\sqrt{(a^2 - b^2)}}{a}, \quad e' = \frac{\sqrt{(b^2 - c^2)}}{b}. \quad 54. \quad \text{A sphere.}$$

55. The polar equation to the locus is

$$a^2 b^2 = (r^2 + c^2) (a^2 \sin^2 \theta + b^2 \cos^2 \theta).$$

$$56. \quad \text{The straight line } \frac{x}{l} = \frac{y}{m} = \frac{z}{n}.$$

58. The given equation may be written thus,  $r^2 = (l^2 + m^2 + n^2) p^2$  where  $r$  is the distance of the point  $(x, y, z)$  from the origin, and  $p$  is the perpendicular from the origin on the plane  $lx + my + nz = 0$ . If  $l^2 + m^2 + n^2$  is greater than unity the locus is a right circular



cone; the cosine of the semivertical angle is  $\frac{1}{\sqrt{l^2 + m^2 + n^2}}$ , If  $l^2 + m^2 + n^2$  is equal to unity the locus is a right line; see example 12. If  $l^2 + m^2 + n^2$  is less than unity the locus is a point, namely the origin. 59. The locus is a sphere if  $C$  be an acute angle, a point if  $C$  be a right angle, and impossible if  $C$  be an obtuse angle.

$$60. \quad k^2 = \frac{a^2}{a^2 - b^2}, \quad m^2 = \frac{a^2(b^2 - c^2)}{c^2(a^2 - b^2)}.$$

62. The equations to the cones are

$$\frac{(x-a)^2}{a^2} = \frac{y^2}{b^2} + \frac{z^2}{c^2}, \quad \frac{(y-b)^2}{b^2} = \frac{z^2}{c^2} + \frac{x^2}{a^2}.$$

63. Let  $(\alpha, \beta, \gamma)$  be the external point; then the required equation is

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = \left( \frac{x\alpha}{a^2} + \frac{y\beta}{b^2} + \frac{z\gamma}{c^2} \right)^2.$$

66. Let  $(\alpha, \beta, \gamma)$  be the given point; then the required equation is

$$a^2(b^2 - c^2)\frac{x}{a} + b^2(c^2 - a^2)\frac{y}{\beta} + c^2(a^2 - b^2)\frac{z}{\gamma} = 0.$$

67. A right circular cone. 70. Take for the equation to the ellipsoid  $\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$ , and for the equation to the sphere  $(x-h)^2 + y^2 + z^2 = r^2$ ; then the equation to the cylinder will be

$$(x-h)^2 + y^2 - \frac{c^2x^2}{a^2} - \frac{c^2y^2}{b^2} = r^2 - c^2.$$

71. The circle circumscribed about the opposite face may be determined by the equations

$$\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1, \quad x^2 + y^2 + z^2 = ax + by + cz.$$

A *subcontrary* section is a circular section. The equation to the cone may be written thus

$$(ax + by + cz - k) \left( \frac{x}{a} + \frac{y}{b} + \frac{z}{c} \right) = x^2 + y^2 + z^2 - k \left( \frac{x}{a} + \frac{y}{b} + \frac{z}{c} \right);$$

hence we see that the plane  $ax + by + cz - k = 0$ , cuts the cone in a circle. If the vertex of the cone be at the angular point which is at the other end of the edge  $c$ , the equation to the cone is

$$x^2 + y^2 = (ax + by) \left(1 - \frac{z}{c}\right);$$

and the plane  $ax + by - cz = 0$  is parallel to the subcontrary sections.

74. An hyperbolic paraboloid.

75. The locus is determined by the equation to the ellipsoid combined with  $xyz = \text{constant}$ .

76. Suppose the fixed point in the axis of  $z$ ; then we have to make  $xy$  a maximum while  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = a$  constant.

85. Circles.

88. Let  $\frac{x' - x}{l} = \frac{y' - y}{m} = \frac{z' - z}{n} = r$  be the equations to a line which passes through the point  $(x, y, z)$  and coincides with the given surface. Substitute the values of  $x', y', z'$  in the given equation; we thus obtain a quadratic in  $r$  which ought to be *identically* true. This leads to

$$\frac{l^2}{a^2} + \frac{m^2}{b^2} - \frac{n^2}{c^2} = 0, \quad \frac{lx}{a^2} + \frac{my}{b^2} - \frac{nz}{c^2} = 0.$$

Eliminate  $m$  from these two equations; thus we obtain a quadratic in  $\frac{l}{n}$ , and the product of the roots becomes known. The result may be written thus

$$\frac{l_1 l_2}{n_1 n_2} = \frac{x^2 - a^2}{z^2 + c^2}.$$

$$\text{Similarly } \frac{m_1 m_2}{n_1 n_2} = \frac{y^2 - b^2}{z^2 + c^2}.$$

$$\text{Hence } 1 + \frac{l_1 l_2}{n_1 n_2} + \frac{m_1 m_2}{n_1 n_2} = \frac{z^2 + c^2 + x^2 - a^2 + y^2 - b^2}{z^2 + c^2}.$$

92. The condition is  $amn + bnl + clm = 0$ .

93. When the given conditions hold, the equation becomes

$$(b'c'x + c'a'y + a'b'z)^2 + 2a'b'c'(a''x + b''y + c''z) - a'b'c'f = 0;$$

this represents a parabolic cylinder.

94. Either  $b' = 0$  and  $c'^2 = (a - c)(b - c)$ ,  
or  $c' = 0$  and  $b'^2 = (a - b)(c - b)$ .

96. Take the equation to an enveloping cylinder

$$\left(\frac{l^2}{a^2} + \frac{m^2}{b^2} + \frac{n^2}{c^2}\right) \left(\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} - 1\right) = \left(\frac{lx}{a^2} + \frac{my}{b^2} + \frac{nz}{c^2}\right)^2,$$

and apply the tests in order that it may be a surface of revolution.

97. An hyperbola. 102. If an ellipsoid be cut by a plane through its centre whose direction cosines are  $l, m, n$ , the area of the section is known to be  $\frac{\pi abc}{\sqrt{(l^2 a^2 + m^2 b^2 + n^2 c^2)}}$ . Now if we seek the maximum and minimum values of  $l^2 a^2 + m^2 b^2 + n^2 c^2$ , with the condition that the plane is to contain the line  $\frac{x}{\lambda} = \frac{y}{\mu} = \frac{z}{\nu}$ , we obtain this quadratic

$$\frac{\lambda^2}{u^2 - a^2} + \frac{\mu^2}{u^2 - b^2} + \frac{\nu^2}{u^2 - c^2} = 0;$$

thus the product of the maximum and minimum values is

$$a^2 b^2 c^2 \left(\frac{\lambda^2}{a^2} + \frac{\mu^2}{b^2} + \frac{\nu^2}{c^2}\right).$$

103. The process will resemble that given in Gregory's *Solid Geometry* for finding the area of the section of the ellipsoid by the plane  $lx + my + nz = 0$ . If  $a, \beta, \gamma$  be co-ordinates of the centre of the section it will be found that

$$\frac{a}{a^2 l} = \frac{\beta}{b^2 m} = \frac{\gamma}{c^2 n} = \frac{\delta}{a^2 l^2 + b^2 m^2 + c^2 n^2}.$$

The final result is

$$\frac{(a^2 l^2 + b^2 m^2 + c^2 n^2 - \delta^2) \pi abc}{(a^2 l^2 + b^2 m^2 + c^2 n^2)^{\frac{3}{2}}}.$$

104. The first part may be proved by taking the general equation

$$ax^2 + by^2 + cz^2 + 2a'yz + 2b'zx + 2c'xy + 2a''x + 2b''y + 2c''z + f = 0;$$

and the cutting planes may be supposed parallel to that of  $(x, y)$ . For the latter part we must find the ratio of similarly situated

lines in the two sections, and the square of this ratio will be the ratio of the areas.

105. This is a mode of expressing the result of the two preceding examples. The result may also be obtained very easily by referring the ellipsoid to conjugate diameters, two of which are in a plane parallel to the planes considered.

106. The equation to the ellipsoid is

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = \frac{\alpha x}{a^2} + \frac{y\beta}{b^2} + \frac{z\gamma}{c^2},$$

where  $(\alpha, \beta, \gamma)$  is the external point.

109. Hyperboloid of one sheet.

110. Hyperboloid of one sheet.

111. Hyperbolic paraboloid.

112. Parabolic cylinder.  
if  $f$  be positive.

113. Hyperboloid of two sheets

114. Right circular cylinder.

115. Hyperboloid of revolution of two sheets if  $f$  be positive.

116. Hyperboloid of revolution of two sheets.

117.  $\frac{2c\gamma}{3\beta}$ ,  $\frac{2c\gamma}{3\alpha}$ , 0 are the co-ordinates.

118. There is a line of centres, given by the equations

$$k = 0, \quad h - l + 1 = 0.$$

119.  $-\frac{1}{2}$ , 0, 0 are the co-ordinates.

122. A surface of revolution, the axis of which is  $x = y = z$ .

124. Use the equation in Example 65; the locus is a sphere, the radius of which is  $\sqrt{(a^2 + b^2 + c^2 + h^2 + k^2)}$ .

133. The sine of the angle is  $\frac{a^2 - c^2}{a^2 + c^2}$  supposing  $a, b, c$  in descending order of magnitude.

142. See a similar example in Geometry of Two Dimensions, *Plane Co-ordinate Geometry*, Art. 286.

144. Refer the ellipsoid to conjugate diameters such that the plane containing two of them is parallel to the cutting plane.

157.  $b^2 - ac$  must be positive and  $(b'^2 - ac)(a'^2 - bc) = (a'b' - cc')^2$ .

159. Let  $x \cos \theta_1 + y \cos \theta_2 + z \cos \theta_3 = u$ ,

$$x \sin \theta_1 + y \sin \theta_2 + z \sin \theta_3 = v;$$

then the second equation is  $uv=0$ , and the first is  $(u+v)(u-v)=0$ .

The line of intersection will be found to be determined by

$$\frac{x}{\sin(\theta_2 - \theta_1)} = \frac{y}{\sin(\theta_1 - \theta_3)} = \frac{z}{\sin(\theta_3 - \theta_1)}.$$

160. The other lines are  $x=0$ ,  $y^2 - z^2 = 0$ ;  $y=0$ ,  $z^2 - x^2 = 0$ ;  
 $z=0$ ,  $x^2 - y^2 = 0$ .

161. The cosine of the inclination of the plane to the axis.

182.  $\phi(x, y, z) = \phi(a + x - a, \beta + y - \beta, \gamma + z - \gamma)$

$$= u + (x-a) \frac{du}{da} + (y-\beta) \frac{du}{d\beta} + (z-\gamma) \frac{du}{d\gamma} \\ + \frac{(x-a)^2}{2} \frac{d^2u}{da^2} + \frac{(y-\beta)^2}{2} \frac{d^2u}{d\beta^2} + \frac{(z-\gamma)^2}{2} \frac{d^2u}{d\gamma^2}$$

$$+ (y-\beta)(z-\gamma) \frac{d^2u}{d\beta d\gamma} + (z-\gamma)(x-a) \frac{d^2u}{d\gamma da} + (x-a)(y-\beta) \frac{d^2u}{da d\beta}.$$

The equation to the tangent plane to the surface  $\phi(x, y, z) = 0$  at the point  $(x, y, z)$  is

$$(x' - x) \frac{d\phi}{dx} + (y' - y) \frac{d\phi}{dy} + (z' - z) \frac{d\phi}{dz} = 0;$$

this may be written

$$(x' - x) \left\{ \frac{du}{da} + (x-a) \frac{d^2u}{da^2} + (z-\gamma) \frac{d^2u}{d\gamma da} + (y-\beta) \frac{d^2u}{da d\beta} \right\} + \&c. = 0.$$

Now suppose this plane to pass through the point  $(a, \beta, \gamma)$ , and we obtain a relation which by means of the equation  $\phi(x, y, z) = 0$ , transformed as above, reduces to

$$2u + (x-a) \frac{du}{da} + (y-\beta) \frac{du}{d\beta} + (z-\gamma) \frac{du}{d\gamma} = 0.$$

183. Let a line pass through the point  $(a, \beta, \gamma)$  and through the point  $(x, y, z)$  on the surface; and let  $(x', y', z')$  be any other point on this line; then we have

$$\frac{x-a}{x'-a} = \frac{y-\beta}{y'-\beta} = \frac{z-\gamma}{z'-\gamma} = r \text{ say; therefore}$$

$$x = a + r(x' - a), \quad y = \beta + r(y' - \beta), \quad z = \gamma + r(z' - \gamma).$$

Substitute these values of  $x, y, z$  in  $\phi(x, y, z) = 0$ ; we thus obtain a *quadratic* in  $r$  corresponding to the two points in which the line through  $(a, \beta, \gamma)$  and  $(x', y', z')$  meets the surface. The condition that this quadratic should have *equal* roots leads to the equation to the required cone,  $x', y', z'$  being the variable co-ordinates.

$$185. \quad a'yz + b'zx + c'xy = 0.$$

$$195. \quad \frac{a^4x^2}{a^2 - k^2} + \frac{c^4y^2}{b^2 - k^2} + \frac{c^4z^2}{c^2 - k^2} = a^2x^2 + b^2y^2 + c^2z^2.$$

196. The surface is determined by the equation

$$y^2 - 2ay + \frac{z^2 - 2kxz}{k^2} + a^2 = 0,$$

where  $k$  is a known constant, and  $a = x \left\{ 1 + \sqrt{\left(\frac{2x}{kc}\right) + \frac{x}{kc}} \right\}$ .

$$200. \quad \frac{x^2}{a^2} + \frac{y^2}{b^2} = \left(1 - \frac{z^2}{c^2}\right) \left(\frac{x}{a} + \frac{y}{b}\right)^2.$$

$$201. \quad \left(\frac{x}{a} \pm \frac{y}{b}\right)^2 + \frac{z^2}{c^2} = 1.$$

205. Take for the equations to the lines those in example 34; let  $a$  be the radius of the circle; the required equation is

$$(mxc^2 + cyz)^2 + m^2(mcxz + c^2y)^2 = m^2a^2(c^2 - z^2)^2.$$

212. (1) is not a developable surface; (2) is.

$$214. \quad z + \sqrt{(x^2 + y^2 + z^2)} = \phi\left(\frac{y}{x}\right).$$

$$215. \quad z^{n+1} - k^{n+1} = x^{n+1} \phi\left(\frac{y}{x}\right).$$

$$216. \quad \frac{x^2 + y^2}{a^2} = \frac{(z - c)^2}{c^2}.$$

$$218. \quad (ax + by + cz)^2 = r^2 \{(x + a)^2 + (y + b)^2 + (z + c)^2\}.$$

219. Let the equations to be parabolas be

$$\left. \begin{array}{l} y = 0 \\ z^2 = 4ax \end{array} \right\}, \quad \text{and} \quad \left. \begin{array}{l} z = 0 \\ y^2 = 4a'x \end{array} \right\};$$

the equation to the surface is  $4x - \left\{ \frac{y}{\sqrt{a'}} + \frac{z}{\sqrt{a}} \right\}^2 = 0$ .

225.  $(x^2 + \mu^2 y^2)^2 = \frac{k^2}{z^2}$ , where  $\mu$  and  $k$  are known constants.

227. Use conjugate diameters as axes, two of them being parallel to the plane of the circles, and the third passing through the centres of the circular sections.

230.  $y = \pm a \cos \beta$ .      231.  $\frac{x'x}{a^2} + \frac{y'y}{b^2} = 1$ ,  $z = \frac{c}{\sqrt{2}}$ .

232.  $x' - x = \frac{2y}{a - 2x}(y' - y) = -\frac{2z}{a}(z' - z)$ .

233. The curve is determined by the given equation combined with  $4z^2 = x^2 + y^2$ .

239. They are proportional to  $\frac{d^2x}{ds^2}$ ,  $\frac{d^2y}{ds^2}$ , and  $\frac{d^2z}{ds^2}$ .

240.  $c(x'y - y'x) + ab(z' - z) = 0$ .      241. A plane passing through the line of intersection of the normal plane at the point  $(x, y, z)$  and the consecutive normal plane, and perpendicular to the first normal plane.

$$243. \frac{(h^2 a^2 + k^2 z^2)^{\frac{3}{2}}}{a^2 h^2 \sqrt{(h^2 + k^2)}}$$

245. A circle.

246. A conic section.

251. The curves of greatest inclination to the plane of  $(x, y)$  are determined by the equation to the given surface combined with  $x^2 - y^2 = \text{constant}$ .

252. The curves are determined by combining the given equation with  $x^2 - y^2 = \text{constant}$ .

253.  $x^2 + 3ay = 0$ ,  $27acz + x^3 = 0$ .      257.  $27a(x+y)^2 = 16(z-a)^2$ .

269.  $z = A \tan^{-1} \frac{y}{x} + B$ .      270. The surface is that formed

by the revolution of the catenary  $y = \frac{c}{2}(e^{\frac{z}{c}} + e^{-\frac{z}{c}})$  round the axis of  $x$ .

299. The condition is  $abc + 2a'b'c' - aa'^2 - bb'^2 - cc'^2 = 0$ .

302.  $Aa + B\beta + C\gamma = 0$ , is the equation to the plane which is parallel to the face of which the area is  $D$ , and which passes through the opposite vertex.

305. The equation represents a cone which touches the planes represented by  $u_1 = 0$ ,  $u_2 = 0$ ,  $u_3 = 0$ .

306. The equation represents a cone containing the lines of intersection of the planes  $u_1 = 0$ ,  $u_2 = 0$ ,  $u_3 = 0$ .

313. An ellipsoid, a point, or an impossible locus according as  $a$  is  $>$  or  $<$  0.

314. An hyperboloid of one sheet, a cone, or an hyperboloid of two sheets according as  $a$  is  $>$  or  $<$  0.

315. An elliptic cylinder, a straight line, or an impossible locus according as  $a$  is  $>$  or  $<$  0.

316. An hyperbolic cylinder, two planes, or an hyperbolic cylinder according as  $a$  is  $>$  or  $<$  0.

317. Two parallel planes, one plane, or an impossible locus according as  $a$  is  $>$  or  $<$  0.

318. An elliptic paraboloid. 319. An hyperbolic paraboloid. 320. A parabolic cylinder. See for the last eight questions *The Mathematician*, Vol. III. p. 195.

In examples 321—329 the symbols  $r$ ,  $\theta$ ,  $\phi$  are the usual polar co-ordinates. 321. A right circular cone having its vertex at the origin and its axis coincident with the axis of  $z$ . 322. A plane containing the axis of  $z$ . 323. A sphere having its centre at the origin. 324. A series of right circular cones having their vertices at the origin and their axes coincident with the axis of  $z$ . 325. A series of planes containing the axis of  $z$ . 326. A series of spheres having the origin for centre. 327. A surface of revolution round the axis of  $z$ . 328. A surface such that any section made by a plane which contains the axis of  $z$  is a circle with the origin for centre. 329. A conical surface generated by straight lines which all pass through the origin.

331. A sphere having the origin on its surface.

$$337. \pm abc \left( \frac{x}{a} + \frac{y}{b} + \frac{z}{c} - 1 \right).$$



340. Let  $c$  be a side of the hexagon,  $a$  an edge of the cube; then  $a = \frac{c\sqrt{3}}{\sqrt{2}}$ .

342. Let  $a$  be the edge of the cube; the height of the luminous point above the given plane is  $a\sqrt{2}(1 + \sqrt{2})$ .

347. The figure formed by the revolution of an hyperbola round its conjugate axis. 354. The equation to the surface is

$$2cx^2 + z\{(x+y)^2 - 2c^2\} - 4cxy = 0; \text{ the volume is } \frac{83\pi c^3}{240\sqrt{2}}.$$

380.  $(x^2 + y^2 + z^2)^3 = 6c^3xyz$  where  $c^3$  is the constant volume.

383.  $c^3xyz = a^3(x^2 + y^2)$ . 389. The volume is  $\frac{\pi a^3}{15}$ .

$$392. \left(\frac{x}{a} + \frac{y}{b} + \frac{z}{c} - 1\right)^2 = 2\left(\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} - 1\right).$$

405. The longest axis of the ellipsoid must be inclined to the plane at an angle whose cosine is  $\left(\frac{b^2 - c^2}{a^2 - c^2}\right)^{\frac{1}{2}}$ .

$$406. y^2 \{\phi(z)\}^2 = \{x - \phi(z)\}^2 [c^2 - \{\phi(z)\}^2].$$

Cambridge.

## LIST OF BOOKS

PUBLISHED BY

**MACMILLAN AND CO.**

**ABD-EL-KADER—The Liberation of Abd-el-Kader.**

An Ode. 8vo. sewed. 1s.

**ABDY.—A Historical Sketch of Civil Procedure among the**

Romans. By J. T. ABDY, LL.D., Regius Professor of Civil Law in the University of Cambridge. Crown 8vo. cloth, 4s. 6d.

**ÆSCHYLI Eumenides.**

The Greek Text with English Notes, and an Introduction, containing an Analysis of Müller's Dissertations. By BERNARD DRAKE, M.A., late Fellow of King's College, Cambridge. 8vo. cloth, 7s. 6d.

**ARISTOTLE on the Vital Principle.**

Translated, with Notes. By CHARLES COLLIER, M.D., F.R.S., Fellow of the Royal College of Physicians. Crown 8vo. cloth, 8s. 6d.

**BEAMONT.—Catherine, the Egyptian Slave in 1852. A Tale**

of Eastern Life. By W. J. BEAMONT, M.A., Fellow of Trinity College, Cambridge, late Principal of the English College, Jerusalem. Fcap. 8vo. 5s. 6d.

**BIRKS.—The Difficulties of Belief in connexion with the**

Creation and the Fall. By THOMAS RAWSON BIRKS, M.A., Rector of Kelshall, and EXAMINING CHAPLAIN TO THE LORD BISHOP OF CARLISLE; Author of "The Life of the Rev. E. Bickersteth." Crown 8vo. cloth, 4s. 6d.

**BOLTON.—The Evidences of Christianity.**

By W. J. BOLTON, of Caius College, Cambridge. 8vo. 6s.

**BOOLE.—The Mathematical Analysis of Logic.**

By GEORGE BOOLE, LL.D. Professor of Mathematics, Queen's College, Cork. 8vo. sewed, 5s.

**BRAVE WORDS for BRAVE SOLDIERS and SAILORS.**

Tenth Thousand. 16mo. sewed, 2d.; or 10s. per 100.

[Printed for Distribution.]

**BUTLER (Archer).—Sermons, Doctrinal and Practical.**

By the Rev. WILLIAM ARCHER BUTLER, M.A. late Professor of Moral Philosophy in the University of Dublin. Edited, with a Memoir of the Author's Life, by the Very Rev. THOMAS WOODWARD, M.A. Dean of Down. With Portrait. Fourth Edition. 8vo. cloth, 12s.

No. 2.]

A

**BUTLER (Archer).—A Second Series of Sermons.**

Edited by J. A. JEREMIE, D.D. Regius Professor of Divinity in the University of Cambridge. **Second Edition.** 8vo. cloth, 10s. 6d.

**BUTLER (Archer).—History of Ancient Philosophy.**

A Series of Lectures. Edited by WILLIAM HEPWORTH THOMPSON, M.A. Regius Professor of Greek in the University of Cambridge. 2 vols. 8vo. cloth, 1l. 5s.

**BUTLER (Archer).—Letters on Romanism, in Reply to Mr.**

NEWMAN's Essay on Development. Edited by the Very Rev. T. WOODWARD, Dean of Down. 8vo. cloth, 10s. 6d.

**CAMBRIDGE.—Cambridge Theological Papers. Comprising**

those given at the Voluntary Theological and Crosse Scholarship Examinations. Edited, with References and Indices, by A. P. MOOR, M.A. of Trinity College, Cambridge, and Sub-warden of St. Augustine's College, Canterbury. 8vo. cloth, 7s. 6d.

**CAMBRIDGE SENATE-HOUSE PROBLEMS and RIDERS, with SOLUTIONS:—**

1848—1851.—Problems. By N. M. FERRERS, M.A. and J. S. JACKSON, M.A. of Caius College. 15s. 6d.

1848—1851.—Riders. By F. J. JAMESON, M.A. of Caius College. 7s. 6d.

1854—Problems and Riders. By W. WALTON, M.A. of Trinity College, and C. F. MACKENZIE, M.A. of Caius College. 10s. 6d.

1857—Problems and Riders. By W. M. CAMPION, M.A. of Queen's College, and W. WALTON, M.A. of Trinity College. 8s. 6d.

**CAMBRIDGE FITZWILLIAM MUSEUM.—A Hand-Book**

to the Pictures in the Fitzwilliam Museum, Cambridge. Crown 8vo. sewed, 1s. 6d.; or in cloth elegant, 2s. 6d.

**CAMBRIDGE.—Cambridge Mathematical Journal. Vol. I.**

Second Edition, 8vo. cloth, 18s.

**CAMBRIDGE.—Cambridge and Dublin Mathematical Journal.**

*The Complete Work*, in Nine Vols. 8vo. cloth, 7l. 4s.

ONLY A FEW COPIES OF THE COMPLETE WORK REMAIN ON HAND.

**CAMPBELL.—The Nature of the Atonement and its Relation**

to Remission of Sins and Eternal Life. By JOHN M'LEOD CAMPBELL, formerly Minister of Row. 8vo. cloth, 10s. 6d.

**CICERO.—Old Age and Friendship.**

Translated into English. Two Parts. 12mo. sewed, 2s. 6d. each.

**COLENZO.—The Colony of Natal. A Journal of Ten Weeks'**

Tour of Visitation among the Colonists and Zulu Kafirs of Natal. By the Right Rev. JOHN WILLIAM COLENZO, D.D. Lord Bishop of Natal, with a Map and Illustrations. Fcap. 8vo. cloth, 5s.

**COLENZO.—Village Sermons.**

Second Edition. Fcap. 8vo. cloth, 2s. 6d.

**COLENZO.—Four Sermons on Ordination, and on Missions.**

18mo. sewed, 1s.

**COLENZO.—Companion to the Holy Communion, containing**

the Service, and Select Readings from the writings of Mr. MAURICE, Edited by the Lord Bishop of Natal. *Fine Edition*, rubricated and bound in morocco, antique style, 6s.; or in cloth, 2s. 6d. *Common Paper*, limp cloth, 1s.

**COTTON.—Sermons: chiefly connected with Public Events**

of 1854. By G. E. LYNCH COTTON, M.A. Master of Marlborough College, formerly Fellow of Trinity College, Cambridge. Fcap. 8vo. cloth, 3s.

**CROSSE.—An Analysis of Paley's Evidences.**

By C. H. CROSSE, M.A. of Caius College, Cambridge. 12mo. boards, 3s. 6d.

**DAVIES.—St. Paul and Modern Thought:**

Remarks on some of the Views advanced in Professor Jowett's Commentary on St. Paul. By Rev. J. LL. DAVIES, M.A. Fellow of Trinity College, Cambridge, and Rector of Christ Church, Marylebone. 8vo. sewed, 2s. 6d.

**DEMOSTHENES.—Demosthenes de Corona.**

The Greek Text, with English Notes. By BERNARD DRAKE, M.A. late Fellow of King's Coll. Cambridge, Editor and Translator of the "Eumenides of Æschylus." Crown 8vo. cloth, 5s.

**DEMOSTHENES.—Demosthenes on the Crown.**

Translated by J. F. NORRIS, M.A. Fellow of Trinity College, Cambridge, and one of Her Majesty's Inspectors of Schools. Crown 8vo. cloth, 3s.

**DRAKE.—Notes, Critical and Explanatory, on Jonah and**

Hosea. By WILLIAM DRAKE, M.A. formerly Fellow of St. John's College, Cambridge. 8vo. cloth, 9s.

**DREW.—A Geometrical Treatise on Conic Sections, with**

Copious Examples from the Cambridge Senate House Papers. By W. H. DREW, M.A. of St. John's College, Cambridge, Second Master of Blackheath Proprietary School. Crown 8vo. cloth, 4s. 6d.

**EVANS.—Sonnets on the Death of Wellington.**

By SEBASTIAN EVANS, of Emmanuel College, Cambridge. 8vo. sewed, 1s.

**FARRAR.—The Influence of Classical Studies on English**

Literature in the Reigns of Elizabeth and James I. By F. W. FARRAR, Fellow of Trinity College, Cambridge, and Assistant Master of Harrow School. 8vo. sewed, 1s. 6d.

**FARRAR.—The Truth shall make you Free.**

A Sermon preached on April 26, 1857. Crown 8vo. sewed, 6d.

**FLOWERS OF THE FOREST. A War Ballad for 1855.**

By M. E. H. Crown 8vo. sewed, 6d.

**FROST.—The First Three Sections of Newton's Principia.**

With Notes and Problems in illustration of the subject. By PERCIVAL FROST, M.A. late Fellow of St. John's College, Cambridge, and Mathematical Lecturer of Jesus College. Crown 8vo. cloth, 10s. 6d.

**GILL.—The Anniversaries. Poems in Commemoration of**

Great Men and Great Events. By T. H. GILL. Fcap. 8vo. cloth, 5s.

**GODFRAY.—An Elementary Treatise on the Lunar Theory.**

With a brief Sketch of the History of the Problem up to the time of Newton. By HUGH GODFRAY, M.A. of St. John's College, Esquire Bedell in the University of Cambridge. 8vo. cloth, 5s. 6d.

**GRANT.—Plane Astronomy.**

Including Explanations of Celestial Phenomena, and Descriptions of Astronomical Instruments. By A. R. GRANT, M.A., one of Her Majesty's Inspectors of Schools, late Fellow of Trinity College, Cambridge. 8vo. boards, 6s.

**GRIFFIN.—The Theory of Double Refraction.**

By W. N. GRIFFIN, B.D. Vicar of Ospringe, Faversham. 8vo. sewed, 2s.

**HALE.—The Transportation Question; or, Why Western**

Australia should be made a Reformatory Colony instead of a Penal Settlement. By the Right Rev. MATTHEW B. HALE, D.D. Lord Bishop of Perth. Crown 8vo. sewed 2s. 6d.

**HALLIFAX.—Analysis of the Civil Law, being a comparison**

of the Roman and English Law. By BISHOP HALLIFAX, with Additions, by J. W. GELDART, LL.D. of Trinity Hall, Cambridge. 8vo. boards, 8s. 6d.

**HAMILTON.—On Truth and Error: Thoughts, in Prose and**

Verse, on the Principles of Truth, and the Causes and Effects of Error. By JOHN HAMILTON, Esq. (of St. Ernan's), M.A. St. John's College, Cambridge. Crown 8vo. cloth, 10s. 6d.

**HARE.—Charges delivered during the Years 1840 to 1854.**

With Notes on the Principal Events affecting the Church during that period. By JULIUS CHARLES HARE, M.A. sometime Archdeacon of Lewes, and Chaplain in Ordinary to the Queen. With an Introduction, explanatory of his position in the Church with reference to the parties which divide it, 3 vols. 8vo. cloth, 1l. 11s. 6d.

**HARE.—Miscellaneous Pamphlets on some of the Leading**

Questions agitated in the Church during the Years 1845—51. 8vo. cloth; 12s.

**HARE.—The Victory of Faith.**

Second Edition. 8vo. cloth, 5s.

**HARE.—The Mission of the Comforter.**

Second Edition. With Notes. 8vo. cloth, 12s.

**HARE.—Vindication of Luther from his English Assailants.**

Second Edition. 8vo. cloth, 7s.

**HARE.—Parish Sermons.**

Second Series. 8vo. cloth, 12s.

**HARE.—Sermons Preacht on Particular Occasions.**

8vo. cloth, 12s.

\*.\* The two following Books are included in the Three Volumes of Charges, and may still be had separately.

**HARE.—The Contest with Rome.**

With Notes, especially in answer to Dr. Newman's Lectures on Present Position of Catholics. Second Edition. 8vo. cloth, 10s. 6d.

**HARE.—Charges delivered in the Years 1843, 1845, 1846.**

Never before published. With an Introduction, explanatory of his position in the Church with reference to the parties which divide it. 6s. 6d.

**HARE.—Portions of the Psalms in English Verse.**

Selected for Public Worship. 18mo. cloth, 2s. 6d.

**HARE.—Two Sermons preached in Herstmonceux Church,**

on Septuagesima Sunday, 1855, being the Sunday after the Funeral of the Venerable Archdeacon Hare. By the Rev. H. VENN ELLIOTT, Perpetual Curate of St. Mary's, Brighton, late Fellow of Trinity College, Cambridge, and the Rev. J. N. SIMPKINSON, Rector of Brighton, Northampton, formerly Curate of Herstmonceux. 8vo. 1s. 6d.

**HARDWICK.—Christ and other Masters.**

A Historical Inquiry into some of the chief Parallelisms and Contrasts between Christianity and the Religious Systems of the Ancient World. With special reference to prevailing Difficulties and Objections. By CHARLES HARDWICK, M.A., Christian Advocate in the University of Cambridge. PART I. Introduction. PART II. THE RELIGIONS OF INDIA. PART III. THE RELIGIONS OF CHINA, AMERICA, AND OCEANICA. 8vo. cloth. 7s. 6d. each part.

**HARDWICK.—A History of the Christian Church, from Gregory the Great to the Reformation. (A.D. 590-1600.)**

Two vols. crown 8vo. cloth, 21s.

Vol. I. contains The History from Gregory the Great to the Excommunication of Luther. With Maps.

Vol. II. contains The History of the Reformation in the Church.

Each volume may be had separately. Price 10s. 6d.

\*.\* These Volumes form part of the Series of Theological Manuals.

**HARDWICK.**—Twenty Sermons for Town Congregations.

Crown 8vo. cloth, 6s. 6d.

**HARDWICK.**—The Universities and the Church of England.

8vo. sewed, 6d.

**HEMMING.**—An Elementary Treatise on the Differential and Integral Calculus. By G. W. HEMMING, M.A. Fellow of St. John's College, Cambridge. **Second Edition.** 8vo. cloth, 9s.**HENSLEY.**—A Sermon Preached on Commemoration Day, 1855, in Trinity College Chapel. By L. HENSLEY, M.A. Vicar of Hitchin, Herts. 8vo. 1s.**HERVEY.**—The Genealogies of our Lord and Saviour Jesus Christ, as contained in the Gospels of St. Matthew and St. Luke, reconciled with each other and with the Genealogy of the House of David, from Adam to the close of the Canon of the Old Testament, and shown to be in harmony with the true Chronology of the Times. By Lord ARTHUR HERVEY, M.A. Rector of Ickworth. 8vo. cloth, 10s. 6d.**HERVEY.**—The Inspiration of Holy Scripture.

Five Sermons preached before the University of Cambridge. 8vo. cloth, 3s. 6d.

**HERVEY.**—A Suggestion for supplying the Literary Institutes of England with Lecturers from the Universities. 8vo. sewed, 6d.**HOWARD.**—The Pentateuch; or, the Five Books of Moses.

Translated into English from the Version of the LXX. With Notes on its Omissions and Insertions, and also on the Passages in which it differs from the Authorised Version. By the Hon. HENRY HOWARD, D.D. Dean of Lichfield. Crown 8vo. cloth. Genesis, 1 vol. 8s. 6d.; Exodus and Leviticus, 1 vol. 10s. 6d.; Numbers and Deuteronomy, 1 vol. 10s. 6d.

**HUMPHREYS.**—Exercitationes Iambicæ; or, ProgressiveExercises in Greek Iambic Verse. To which are prefixed the Rules of Greek Prosody, with copious Notes and Illustrations of the Exercises. By E. R. HUMPHREYS, LL.D. Head Master of the Cheltenham Grammar School. **Second Edition.** Fcap. cloth, 5s. 6d.**INGLEBY.**—Outlines of Theoretical Logic.

Founded on the New Analytic of SIR WILLIAM HAMILTON. Designed for a Text-book in Schools and Colleges. By C. MANSFIELD INGLEBY, M.A. of Trinity College, Cambridge. In fcap. 8vo. cloth, 3s. 6d.

**JEWELL.**—An Apology of the Church of England, and an

Epistle to Seignor Scipio concerning the Council of Trent, translated from the original Latin, and illustrated with Notes, chiefly drawn from the Author's "Defence of the Apology." By A. T. RUSSELL. Fcp. 8vo. bds. 5s.

**JUSTIN MARTYR.**—S. *Justini Philosophi et Martyris Apologia Prima.* Edited, with a corrected Text, and English Introduction with explanatory Notes, by W. TROLLOPE, M.A. Pembroke College, Cambridge. 8vo. bds. 7s. 6d.

**JUVENAL.**—Juvenal, for Schools.

With English Notes. By J. E. B. MAYOR, M.A. Fellow and Classical Lecturer of St. John's College, Cambridge. Crown 8vo. cloth, 10s. 6d.

**KENNEDY.**—The Influence of Christianity on International Law. By C. M. KENNEDY, of Caius College, Cambridge. Crown 8vo. cloth, 4s.

**KINGSLEY.**—Two Years Ago.

By CHARLES KINGSLEY, F.S.A. Author of "Hypatia." **Second Edition.** 3 vols. crown 8vo. cloth, 11. 11s. 6d.

**KINGSLEY.**—"Westward Ho!" or, the Voyages and Adventures of Sir Amyas Leigh, Knight of Burrough, in the County of Devon, in the Reign of Her Most Glorious Majesty Queen Elizabeth. **Third Edition.** Crown 8vo. cloth, 7s. 6d.

**KINGSLEY.**—The Heroes: or, Greek Fairy Tales for my Children. With Eight Illustrations by the Author. In 8vo. beautifully printed on tinted paper, and elegantly bound in cloth, with gilt leaves, 7s. 6d.

**KINGSLEY.**—Glaucus; or, the Wonders of the Shore.

**Third Edition,** corrected and enlarged. With a Frontispiece. Fcap. 8vo. elegantly bound in cloth, with gilt leaves, 3s. 6d.

**KINGSLEY.**—Alexandria and Her Schools: being Four Lectures delivered at the Philosophical Institution, Edinburgh. With a Preface. Crown 8vo. cloth, 5s.

**KINGSLEY.**—Phaethon; or Loose Thoughts for Loose Thinkers. **Second Edition.** Crown 8vo. boards, 2s.

**LECTURES to Ladies on Practical Subjects.**

By Rev. F. D. MAURICE.  
Rev. C. KINGSLEY.  
Dr. GEORGE JOHNSON.  
Dr. SIEVEKING.  
Rev. J. LL. DAVIES.  
Dr. CHAMBERS.

By F. J. STEPHEN, Esq.  
Archdeacon ALLEN.  
Dean TRENCH.  
TOM TAYLOR, Esq.  
Rev. J. S. BREWER.

**Third Edition.** 7s. 6d.

**LUDLOW.**—British India; its Races, and its History, considered with reference to the Mutinies of 1857. A Series of Lectures. By JOHN MALCOLM LUDLOW, Barrister-at-Law. 2 vols. fcap. 8vo. cloth, 9s.



**LUSHINGTON.**—*La Nation Boutiquière: and other Poems*, chiefly Political. With a Preface. By the late HENRY LUSHINGTON, Chief Secretary to the Governor of Malta. *Points of War.* By FRANKLIN LUSHINGTON, Judge in the Supreme Courts of the Ionian Isles. In 1 vol. fcap. 8vo. cloth, 3s.

**MACKENZIE.**—*The Christian Clergy of the first Ten Centuries, and their Influence on European Civilization.* By HENRY MACKENZIE. B.A. Scholar of Trinity College, Cambridge. Crown 8vo. cloth, 6s. 6d.

**MACLEAR.**—*Incentives to Virtue, Natural and Revealed.* By G. F. MACLEAR, B.A. Scholar of Trinity College, Cambridge. Crown 8vo. sewed, 1s. 6d.

**MANSFIELD.**—*Paraguay, Brazil, and the Plate.* With a Map, and numerous Woodcuts. By CHARLES MANSFIELD, M.A. of Clare College, Cambridge. With a Sketch of his Life. By the Rev. CHARLES KINGSLEY. Crown 8vo. cloth, 12s. 6d.

**M'COY.**—*Contributions to British Palæontology; or, First Descriptions of several hundred Fossil Radiata, Articulata, Mollusca, and Pisces, from the Tertiary, Cretaceous, Oolitic, and Palæozoic Strata of Great Britain.* With numerous Woodcuts. By FREDERICK M'COY F.G.S., Professor of Natural History in the University of Melbourne. 8vo. cloth, 9s.

**MASSON.**—*Essays, Biographical and Critical; chiefly on the English Poets.* By DAVID MASSON, M.A. Professor of English Literature in University College, London. 8vo. cloth, 12s. 6d.

**MAURICE.**—*Expository Works on the Holy Scriptures:—*

**I.—The Patriarchs and Lawgivers of the Old Testament.**  
**Second Edition.** Crown 8vo. cloth, 6s.

This volume contains Discourses on the Pentateuch, Joshua, Judges, and the beginning of the First Book of Samuel.

**II.—The Prophets and Kings of the Old Testament.**

**Second Edition.** Crown 8vo. cloth, 10s. 6d.

This volume contains Discourses on Samuel I. and II., Kings I. and II., Amos, Joel, Hosea, Isaiah, Micah, Nahum, Habakkuk, Jeremiah, and Ezekiel.

**III.—The Unity of the New Testament.**

8vo. cloth, 14s.

This Volume contains Discourses on the Gospels of St. Matthew, St. Mark, and St. Luke; the Acts of the Apostles; the Epistles of St. James, St. Jude, St. Peter, and St. Paul.

**IV.—The Gospel of St. John; a Series of Discourses.**

**Second Edition.** Crown 8vo. cloth, 10s. 6d.

**V.—The Epistles of St. John; a Series of Lectures on Christian Ethics.** Crown 8vo. cloth, 7s. 6d.

**MAURICE.—Expository Works on the Prayer-Book.**

**I.—The Ordinary Services.**

**Second Edition.** Fcap. 8vo. cloth, 5s. 6d.

**II.—The Church a Family. Twelve Sermons on the**

Occasional Services. Fcap. 8vo. cloth, 4s. 6d.

**MAURICE.—Lectures on Ecclesiastical History.**

By **FREDERICK DENISON MAURICE, M.A.**, Chaplain of Lincoln's Inn.  
8vo. cloth, 10s. 6d.

**MAURICE.—Theological Essays.**

**Second Edition**, with a new Preface and other additions. Crown 8vo.  
cloth, 10s. 6d.

**MAURICE.—The Doctrine of Sacrifice deduced from the**

Scriptures. With a Dedicatory Letter to the Young Men's Christian Association.  
Crown 8vo. cloth, 7s. 6d.

**MAURICE.—Christmas Day, and other Sermons.**

8vo. cloth, 10s. 6d.

**MAURICE.—The Religions of the World, and their Relations**

to Christianity. **Third Edition.** Fcap. 8vo. cloth, 5s.

**MAURICE.—On the Lord's Prayer.**

**Third Edition.** Fcap. 8vo. cloth, 2s. 6d.

**MAURICE.—On the Sabbath Day: the Character of the**

Warrior; and on the Interpretation of History. Fcap. 8vo. cloth, 2s. 6d.

**MAURICE.—Learning and Working.—Six Lectures on the**

Foundation of Colleges for Working Men, delivered in Willis's Rooms,  
London, in June and July, 1854. Crown 8vo. cloth, 5s.

**MAURICE.—The Indian Crisis. Five Sermons.**

Crown 8vo. cloth, 2s.

**MAURICE.—Law's Remarks on the Fable of the Bees.**

Edited, with an Introduction of Eighty Pages, by **FREDERICK DENISON MAURICE, M.A.** Chaplain of Lincoln's Inn. Fcp. 8vo. cloth, 4s. 6d.

**MAURICE.—Miscellaneous Pamphlets:—**

**I.—Eternal Life and Eternal Death.**

Crown 8vo. sewed, 1s. 6d.

**II.—Death and Life. A Sermon. In Memoriam C. E. G.**

8vo. sewed, 1s.

**MAURICE.**—Miscellaneous Pamphlets, continued:—

**III.**—Plan of a Female College for the Help of the Rich and of the Poor. 8vo. 6d.

**IV.**—Administrative Reform.  
Crown 8vo. 3d.

**V.**—The Word "Eternal," and the Punishment of the Wicked. Fifth Thousand. 8vo. 1s.

**VI.**—The Name "Protestant;" and the English Bishopric at Jerusalem. Second Edition. 8vo. 3s.

**VII.**—Right and Wrong Methods of Supporting Protestantism. 8vo. 1s.

**VIII.**—Thoughts on the Oxford Election of 1847.  
8vo. 1s.

**IX.**—The Case of Queen's College, London.  
8vo. 1s. 6d.

**X.**—The Worship of the Church a Witness for the Redemption of the World. 8vo. sewed, 1s.

**MAYOR.**—Cambridge in the Seventeenth Century.

2 vols. fcap. 8vo. cloth, 13s.

Vol. I. contains The Lives of Nicholas Ferrar.

Vol. II. contains The Autobiography of Matthew Robinson.

By JOHN E. B. MAYOR, M.A. Fellow and Assistant Tutor of St. John's College, Cambridge.

\* \* \* The Autobiography of Matthew Robinson may be had separately, price 5s. 6d.

**MINUCIUS FELIX.**—The Octavius of Minucius Felix.

Translated into English by LORD HAILES. Fcap. 8vo. cloth, 3s. 6d.

**MORSE.**—Working for God; Four Sermons Preached before the University of Cambridge. By FRANCIS MORSE, M.A. Incumbent of St. John's, Ladywood, Birmingham. Crown 8vo. sewed, 2s. 6d.

**NAPIER.**—Lord Bacon and Sir Walter Raleigh.

Critical and Biographical Essays. By MACVEY NAPIER, late Editor of the *Edinburgh Review* and of the *Encyclopædia Britannica*. Post 8vo. cloth, 7s. 6d.

**NIND.**—Sonnets of Cambridge Life. By Rev. W. NIND, M.A. late Fellow of St. Peter's College. Post 8vo. boards, 2s.

**NIND.**—The German Lyrist; or, Metrical Versions from the principal German Lyric Poets. Crown 8vo. cloth, 3s.

**NIND.**—The Odes of Klopstock Translated.

Fcap. 8vo. cloth, 6s.

**NIND.**—Italy and Vienna in 1851.

A Series of Letters. Fcap. 8vo. cloth, 5s. 6d.

**NORRIS.**—Ten School-Room Addresses.

Edited by J. P. NORRIS, M.A. Fellow of Trinity College, and one of Her Majesty's Inspectors of Schools. 18mo. sewed, 8d.

**NORWAY AND SWEDEN.**—A Long Vacation Ramble in

1856. By X and Y. Crown 8vo. cloth, 6s. 6d.

**PARKINSON.**—A Treatise on Elementary Mechanics.

For the Use of the Junior Classes at the University, and the Higher Classes in Schools. With a Collection of Examples. By S. PARKINSON, M.A. Fellow and Assistant Tutor of St. John's College, Cambridge. Crown 8vo. cloth, 9s. 6d.

**PARMINTER.**—Materials for a Grammar of the Modern

English Language. Designed as a Text-book of Classical Grammar for the use of Training Colleges, and the Higher Classes of English Schools. By GEORGE HENRY PARMINTER, of Trinity College, Cambridge; Rector of the United Parishes of SS. John and George, Exeter. Fcap. 8vo. cloth, 3s. 6d.

**PAYN.**—Poems.

By JAMES PAYN. Fcp. 8vo. cloth, 5s.

**PAYNE.**—Decaeus; or, the Dawn of To-morrow, and The

Bondchild. Two Sacred Allegories. By ISABEL J. PAYNE. 18mo. cloth, 1s.

**PEACE IN WAR.** *In Memoriam* E. K.

Crown 8vo. sewed, 1s.

**PEROWNE.**—"Al-Adjrumiieh."

An Elementary Arabic Grammar. By J. J. S. PEROWNE, B.D. Fellow of Corpus Christi College, Cambridge, and Lecturer in Divinity in King's College, London. 8vo. cloth, 5s.

**PERRY.**—Sermons Preached before the University of Cam-

bridge. By the Right Rev. CHARLES PERRY, Lord Bishop of Melbourne, formerly Fellow and Tutor of Trinity College, Cambridge. Crown 8vo. cloth, 3s.

**PHEAR.**—Elementary Hydrostatics.

Second Edition. Accompanied by numerous Examples, with the Solutions. Crown 8vo. cloth, 5s. 6d.

**PLAIN RULES ON REGISTRATION OF BIRTHS AND****DEATHS.** Crown 8vo. sewed, 1d.; 9d. per dozen; 5s. per 100.

**PLATO.—The Republic of Plato.**

Translated into English, with Notes. By Two Fellows of Trinity College, Cambridge, (J. Ll. Davies M.A., and D. J. Vaughan, M.A.) **Second Edition.** 8vo. cloth.

**PRINCIPLES of ETHICS according to the NEW TESTAMENT.** Crown 8vo. sewed, 2s.**PROCTER.—A History of the Book of Common Prayer: with a Rationale of its Offices.** By FRANCIS PROCTER, M.A., Vicar of Witton, Norfolk, and late Fellow of St. Catherine's College. **Third Edition,** revised and enlarged. Crown 8vo. cloth, 10s. 6d.

\*.\* This forms part of the Series of Theological Manuals.

**PUCKLE.—An Elementary Treatise on Conic Sections and Algebraic Geometry.** With a numerous collection of Easy Examples progressively arranged, especially designed for the use of Schools and Beginners. By G. HALE PUCKLE, M.A., Principal of Windermere College. **Second Edition,** enlarged and improved. Crown 8vo. cloth, 7s. 6d.**PURCHAS.—The Priest's Dream.**

By JOHN PURCHAS, M.A. of Christ's College, Cambridge. 8vo. sewed, 1s.

**PURTON.—Corporate Life.**

A Sermon Preached before the University of Cambridge. By J. S. PURTON, B.D. Fellow and Tutor of St. Catherine's College. 8vo. sewed, 1s.

**RAMSAY.—The Catechiser's Manual; or, the Church Catechism illustrated and explained, for the use of Clergymen, Schoolmasters, and Teachers.** By ARTHUR RAMSAY, M.A. of Trinity College, Cambridge. 18mo. cloth, 3s. 6d.**REICHEL.—The Lord's Prayer and other Sermons.**

By C. P. REICHEL, B.D., Professor of Latin in the Queen's University; Chaplain to his Excellency the Lord-Lieutenant of Ireland; and late Donnellan Lecturer in the University of Dublin. Crown 8vo. cloth, 7s. 6d.

**ROBINSON.—Missions urged upon the State, on Grounds both of Duty and Policy.** By C. K. ROBINSON, M.A. Fellow and Assistant Tutor of St. Catherine's College. Fcap. 8vo. cloth, 3s.**RUTH AND HER FRIENDS. A Story for Girls.**

With a Frontispiece. Fcap. 8vo. cloth, 5s.

**SALLUST.—Sallust for Schools.**

With English Notes. By CHARLES MERIVALE, B.D.; late Fellow and Tutor of St. John's College, Cambridge, &c., Author of the "History of Rome," &c. Crown 8vo. cloth, 5s.

**SEDGWICK AND M'COY'S British Palæozoic Fossils,**  
Classified and Described. By ADAM SEDGWICK, F.R.S., Professor of Geology  
in the University of Cambridge, and FREDERICK M'COY, F.G.S., Professor of  
Natural History in the University of Melbourne. With numerous Plates.  
Two Vols. 4to. cloth, £2 2s.

**SELWYN.—The Work of Christ in the World.**

Sermons preached before the University of Cambridge. By the Right Rev.  
GEORGE AUGUSTUS SELWYN, D.D. Bishop of New Zealand, formerly  
Fellow of St. John's College. **Third Edition.** Crown 8vo. 2s.

**SELWYN.—A Verbal Analysis of the Holy Bible.**

Intended to facilitate the translation of the Holy Scriptures into Foreign  
Languages. Compiled for the use of the Melanesian Mission. Small folio,  
cloth, 14s.

**SIMPSON.—An Epitome of the History of the Christian**

Church during the first Three Centuries and during the Reformation. With  
Examination Papers. By WILLIAM SIMPSON, M.A. **Third Edition,**  
Fcp. 8vo. cloth, 5s.

**SMITH.—City Poems.**

By ALEXANDER SMITH, Secretary to the University of Edinburgh.  
Author of "A Life Drama," and other Poems. Fcp. 8vo. cloth. 5s.

**SMITH.—Arithmetic and Algebra, in their Principles and**

Application: with numerous systematically arranged Examples, taken from  
the Cambridge Examination Papers. With especial reference to the ordinary  
Examination for B.A. Degree. By BARNARD SMITH, M.A., Fellow of St.  
Peter's College, Cambridge. **Fifth Edition.** Crown 8vo. cloth, 10s. 6d.

**SMITH.—Arithmetic for the use of Schools. By BARNARD**

SMITH, M.A. Fellow of St. Peter's College. **New Edition.** Crown 8vo.  
cloth, 4s. 6d.

**SMITH.—A Key to the Arithmetic for Schools.**

Crown 8vo. cloth, 8s. 6d.

**SNOWBALL.—The Elements of Plane and Spherical**

Trigonometry. By J. C. SNOWBALL, M.A. Fellow of St. John's College,  
Cambridge. **Ninth Edition.** Crown 8vo. cloth, 7s. 6d.

**SNOWBALL.—Introduction to the Elements of Plane Trigo-**

nometry for the use of Schools. **Second Edition.** 8vo. sewed, 5s.

**SNOWBALL.—The Cambridge Course of Elementary**

Mechanics and Hydrostatics. Adapted for the use of Colleges and Schools.  
With numerous Examples and Problems. **Fourth Edition.** Crown 8vo.  
cloth, 5s.

**SWAINSON.—A Handbook to Butler's Analogy.**

By C. A. SWAINSON, M.A. Principal of the Theological College, and  
Prebendary of Chichester. Crown 8vo. sewed, 2s.

**TAIT and STEELE.—A Treatise on Dynamics, with nume-**

rous Examples. By F. G. TAIT, Fellow of St. Peter's College, Cambridge,  
and Professor of Mathematics in Queen's College, Belfast, and W. J. STEELE,  
late Fellow of St. Peter's College. Crown 8vo. cloth, 10s. 6d.

**TAYLOR.—The Restoration of Belief.**

By ISAAC TAYLOR, Esq., Author of "The Natural History of Enthusa-  
siasm." Crown 8vo. cloth, 8s. 6d.

**THEOLOGICAL Manuals.**

CHURCH HISTORY: FROM GREGORY THE GREAT TO THE  
REFORMATION (A.D. 590—1600). By CHARLES HARDWICK. With  
Four Maps, 2 vols. Crown 8vo. cloth, price 1*l.* 1s.

THE COMMON PRAYER: ITS HISTORY AND RATIONALE. By  
FRANCIS PROCTER. Third Edition. Crown 8vo. cloth, 10s. 6d.

A HISTORY OF THE CANON OF THE NEW TESTAMENT. By  
B. F. WESTCOTT. Crown 8vo. cloth, 12s. 6d.

\* \* \* Others are in progress, and will be announced in due time.

**THING.—A Construing Book.**

Compiled by the Rev. EDWARD THING, M.A. Head Master of Up-  
pingham Grammar School, late Fellow of King's College, Cambridge. Fcap.  
8vo. cloth, 2s. 6d.

**THING.—The Elements of Grammar taught in English.**

Second Edition. 18mo. bound in cloth, 2s.

**THING.—The Child's Grammar.**

Being the substance of the above, with Examples for Practice. Adapted for  
Junior Classes. A New Edition. 18mo. limp cloth, 1s.

**THRUPP.—Antient Jerusalem: a New Investigation into the**

History, Topography, and Plan of the City, Environs, and Temple. Designed  
principally to illustrate the records and prophecies of Scripture. With Map  
and Plans. By JOSEPH FRANCIS THRUPP, M.A. Vicar of Barrington,  
Cambridge, late Fellow of Trinity College. 8vo. cloth, 15s.

**THRUPP.—Psalms and Hymns for Public Worship.**

18mo. cloth, 2s.; limp cloth, 1s. 4d.

**THUCYDIDES, BOOK VI. With English Notes, and a Map.**

By PERCIVAL FROST, Jun. M.A. late Fellow of St. John's College,  
Cambridge. 8vo. 7s. 6d.

**TODHUNTER.**—A Treatise on the Differential Calculus; and the Elements of the Integral Calculus. With numerous Examples. By I. TODHUNTER, M.A., Fellow and Assistant Tutor of St. John's College, Cambridge. **Second Edition.** Crown 8vo. cloth, 10s. 6d.

**TODHUNTER.**—A Treatise on the Integral Calculus. With numerous Examples. Crown 8vo. cloth, 10s. 6d.

**TODHUNTER.**—A Treatise on Analytical Statics, with numerous Examples. Crown 8vo. cloth, 10s. 6d.

**TODHUNTER.**—A Treatise on Conic Sections, with numerous Examples. Crown 8vo. cloth, 10s. 6d.

**TOM BROWN'S SCHOOL DAYS.**  
By AN OLD BOY. **Fifth Edition.** Crown 8vo. cloth, 10s. 6d.

**TRENCH.**—Synonyms of the New Testament.  
By The Very Rev. RICHARD CHENEVIX TRENCH, D.D. Dean of Westminster. **Third Edition.** Fcap. 8vo. cloth, 5s.

**TRENCH.**—Hulsean Lectures for 1845—46.  
CONTENTS. 1.—The Fitness of Holy Scripture for unfolding the Spiritual Life of Man. 2.—Christ the Desire of all Nations; or the Unconscious Prophecies of Heathendom. **Third Edition.** Foolscap 8vo. cloth, 5s.

**TRENCH.**—Sermons Preached before the University of Cambridge. Fcap. 8vo. cloth, 2s. 6d.

**VAUGHAN.**—Sermons preached in St. John's Church, Leicester, during the years 1855 and 1856. By DAVID J. VAUGHAN, M.A. Fellow of Trinity College, Cambridge, and Incumbent of St. Mark's, White-chapel. Crown 8vo. cloth, 5s. 6d.

**WAGNER.**—Memoir of the Rev. George Wagner, late of St. Stephen's, Brighton. By J. N. SIMPKINSON, M.A. Rector of Brighton, Northampton. **Second Edition.** Crown 8vo. cloth, 9s.

**WATERS OF COMFORT.**—A Small Volume of Devotional Poetry of a Practical Character. By the Author of "Visiting my Relations." Fcap. 8vo. cloth, 4s.

**WESTCOTT.**—A general View of the History of the Canon of the New Testament during the First Four Centuries. By BROOKE FOSS WESTCOTT, M.A., Assistant Master of Harrow School; late Fellow of Trinity College, Cambridge. Crown 8vo. cloth, 12s. 6d.

\*.\* This forms part of the Series of Theological Manuals.



**WILSON.—The Five Gateways of Knowledge.**

By GEORGE WILSON, M.D., F.R.S.E., Regius Professor of Technology in the University of Edinburgh. **Second Edition.** Fcap. 8vo. cloth, 2s. 6d. or in Paper Covers, 1s.

**WILSON.—A Treatise on Dynamics.**

By W. P. WILSON, M.A., Fellow of St. John's, Cambridge, and Professor of Mathematics in the University of Melbourne. 8vo. bds. 9s. 6d.

**WRIGHT.—Hellenica; or, a History of Greece in Greek,**

as related by Diodorus and Thucydides, being a First Greek Construing Book, with Explanatory Notes, Critical and Historical. By J. WRIGHT, M.A., of Trinity College, Cambridge, and Head-Master of Sutton Coldfield Grammar School. **Second Edition, WITH A VOCABULARY.** 12mo. cloth, 3s. 6d.

**WRIGHT.—A Help to Latin Grammar;**

or, the Form and Use of Words in Latin. With Progressive Exercises. Crown 8vo. cloth, 4s. 6d.

**WRIGHT.—The Seven Kings of Rome:**

An easy Narrative, abridged from the First Book of Livy by the omission of difficult passages, being a First Latin Construing Book, with Grammatical Notes. Fcap. 8vo. cloth, 3s.

**WRIGHT.—A Vocabulary and Exercises on the "Seven**

Kings of Rome." Fcap. 8vo. cloth, 2s. 6d.

---

**THE JOURNAL**

OF

**CLASSICAL AND SACRED PHILOLOGY.**

*No. X. for March 1857, 4s.*

Vols. I. to III. are now ready, 12s. 6d. each, cloth lettered,

CASES CAN BE HAD FOR BINDING THE VOLUMES.

\*.\* Three Numbers published annually, at 4s. each.

---

E. CLAY, PRINTER, BREAD STREET HILL.

